



Use and valuation of native and introduced medicinal plant species in Campo Hermoso and Zetaquirá, Boyacá, Colombia

Cadena-Gonzalez, Ana Lucia; Sørensen, Marten; Theilade, Ida

Published in:
Journal of Ethnobiology and Ethnomedicine

DOI:
[10.1186/1746-4269-9-23](https://doi.org/10.1186/1746-4269-9-23)

Publication date:
2013

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Cadena-Gonzalez, A. L., Sørensen, M., & Theilade, I. (2013). Use and valuation of native and introduced medicinal plant species in Campo Hermoso and Zetaquirá, Boyacá, Colombia. *Journal of Ethnobiology and Ethnomedicine*, 9, [23]. <https://doi.org/10.1186/1746-4269-9-23>



RESEARCH

Open Access

Use and valuation of native and introduced medicinal plant species in Campo Hermoso and Zetaquirá, Boyacá, Colombia

Ana Lucía Cadena-González^{1*}, Marten Sørensen² and Ida Theilade³

Abstract

Background: Medicinal plant species contribute significantly to folk medicine in Colombia. However, few local studies have investigated whether species used are introduced or native and whether there is a difference in importance of native and introduced medicinal plant species. The aim of the present study was to describe the use of medicinal plants within two municipalities, Campo Hermoso and Zetaquirá, both in the department of Boyacá, Colombia and to assess the importance of native and introduced plants to healers, amateur healers and local people. As local healers including amateur healers have no history of introduced species our working hypotheses (H_{1-2}) were that H_1 : native and introduced medicinal plant species are of equal importance and H_2 : healers and amateur healers do not differentiate in their preferences between native and introduced medicinal plant species.

Methods: Ten villages were included in the study. A combination of quantitative and qualitative methods was used including questionnaires, semi-structured interviews, in- depth interviews, and open talks. Voucher specimens were collected in home gardens and during field walks. For data analysis, we calculated use value indices and Jaccard index and tested for the above hypothesis using Spearman rank-correlation coefficients and Wilcoxon-Mann-Whitney tests.

Results: Eighty medicinal plant species were described by locals as the most frequently used. Of these, 78 species were taxonomically identified, distributed within 41 families and 74 genera, which included 35 native species and 43 introduced. The highest valued families were: Asteraceae, Lamiaceae, Apiaceae, Rutaceae and Verbenaceae. The species ranked highest according to their Use Values, in both municipalities, were *Mentha suaveolens* Ehrh., *Ambrosia cumanensis* Kunth, and *Verbena littoralis* Kunth. Introduced species were more important than native ones in Zetaquirá, while there was no difference in importance in Campo Hermoso. While healers relied most on the uses of native species, amateur healers were inclined to rely on introduced species. Medicinal plant administration in both municipalities follow the usual pattern: Leaves are used most commonly prepared by decoction or infusion and administered orally.

Conclusions: The high proportion of introduced plant species used in the local traditional medicines is similar to the results of a number of other ethnobotanical studies and emphasise the need for efforts to record and maintain traditional knowledge on native species.

Keywords: Acculturation, Ethnobotany, Trained healers, Amateur healers, Introduced medicinal plants, Traditional knowledge

* Correspondence: analu@dsr.life.ku.dk

¹Department of Agriculture and Ecology, University of Copenhagen, Rolighedsvej 21, 1958 Frederiksberg C, Denmark

Full list of author information is available at the end of the article

Background

Medicinal plant species constitute important alternatives to conventional medicine in a large number of developing countries, especially within poor communities that inhabit rural areas and lack access to health services [1,2]. A number of native medicinal plants of both the palaeotropics and neotropics have traditionally had a high value for indigenous communities, not only because of their healing properties, but also due to other uses [3]. An example of such multi-use plants is the neotropical species *Crescentia cujete* L., a source of both timber and traditional medicine. Recently, this species has reached the United States and European pharmaceutical markets demonstrating that traditional medicinal plants used by indigenous communities may be of high economic value [3].

However, knowledge, cultivation and maintenance of these native species within rural communities is decreasing, due to modernization processes, such as acculturation [1-3]. In addition, a tendency to consider all plant resources as native by local people has been directly or indirectly documented in several studies [4-6]; in other words, numerous introduced plant species with healing properties have become popularly considered as 'native'. Consequently, a number of native medicinal plant species have been replaced by introduced species. Thus, they use the terms 'native' and 'introduced' indiscriminately so that many introduced species with healing properties have become popularly known as native. For these reasons, the uses of a number of native species have been replaced by introduced species, incorrectly perceived as native.

When faced with the rapid decline in traditional knowledge it is relevant to identify medicinal species and to record their uses within local communities. This is especially important in regions that currently are affected by land-use change and modernisation. Documentation and awareness of ethnobotanical knowledge within these regions may facilitate the maintenance of medicinal plant resources and uses.

Furthermore, ethnobotanical studies may support implementation of strategies that integrate native medicinal plant uses with sustainable management of natural resources. The municipalities of Campo Hermoso and Zetaquirá in the department of Boyacá, Colombia, are examples where conversion of forest areas into agriculture land and reforestation with exotic species have destroyed large areas of natural forest including high plateau, upland, and montane forest. Furthermore, many multiple-use plants have been overexploited [7,8]. This situation has caused loss of a number of native species (pers. comm. local key informants 2009). Studies on medicinal plant use have been conducted in different localities within Boyacá [9,10], but none in the regions of

Campo Hermoso and Zetaquirá. Therefore, the present study was carried out within these municipalities. The aims of this study were to identify the medicinal plant species, to determine their origin (native or introduced to Colombia), to record their medicinal uses (targeted illnesses, ways of administration), and based on personal observations and inspired by previous studies [4-6] to assess the value of native and introduced plant species to members of local communities. The latter objective was evaluated by the null hypotheses (H_{1-2}) H_1 : native and introduced medicinal plant species are of equal importance and H_2 : healers and amateur healers do not differentiate in their preferences between native and introduced medicinal plant species.

Methods

Study areas


The study took place in the municipalities of Campo Hermoso and Zetaquirá in department of Boyacá, Colombia (Figure 1). The municipalities are located on the Eastern slope of the Cordillera Oriental (Eastern Andean range) and their terrain comprises rugged areas with peaks, steep canyons, and valleys. Both study areas belong to the humid subtropical zone [7,8]. The altitude range from 500–2500 m a.s.l. in Campo Hermoso and 1875–3600 m a.s.l. in Zetaquirá. Temperatures range between 12–35°C.

Climatic differences and mountainous topography lead to a diversity of natural vegetation, livestock pastures, and crops within the study areas [7,8]. Campo Hermoso is located at 132 km from Tunja the capital of the department and 143 km from Bogotá. Zetaquirá is located at 69 km away from Tunja, and at 126 km away from Bogotá.

The economy of the municipalities is based on cattle ranching and agriculture; the latter often by smallholders practicing subsistence economy. The main products in Zetaquirá are coffee, beans, sugarcane, maize, plantain, manioc and 'arracacha' (*Arracacia xanthorrhiza* Bancr.) [11]. The coffee production is located near the villages at intermediate altitudes. Until 2008, coffee was a main crop in Zetaquirá. However, climate change has decreased feasibility of coffee production in recent years (pers. comm.: Domingo Mendoza, responsible of the Federación Nacional de Cafeteros de Colombia, office in Zetaquirá). Campo Hermoso is basically a cattle ranching region with large areas of pastures [7]. Besides, maize, beans, manioc and vegetables and, at smaller scale, sugarcane is cultivated for subsistence. There are no conspicuous economic differences between the two municipalities [7,8].

Data collection and free, prior informed consent (FPIC)

The field work was conducted from October 2008 to January 2009. Data collection included a combination of



quantitative and qualitative methods in order to allow for triangulating and cross-checking [12]. Applied and complementary ethnobotanical methods [13] were used to acquire information on how local people related to the medicinal plant resources in the two municipalities. The project was presented to the local authorities and to the communities during initial meetings. Objectives and methods were approved in both municipalities.

Selection of villages and identification of informants

Within each municipality representative villages were visited. Ten villages were chosen according to distance to the main urban centre, the presence of a representative flora, and residential areas of the healers and herbalists, who were purposefully included in the sampling. In Zetaquirá the villages studied were: La Esperanza, Guanata, Centro Rural and Hormigas; and in Campo Hermoso: Papayal, Huerta Vieja, Palmichal, Los Cedros, Macanalito and Teguás.

Participation of local people was essential to several stages of the research process [12]. Thus, key informants

in the municipalities were contacted in the first stage. Their collaboration was crucial for selecting and planning the visits to the villages.

Each municipality was visited three times during the study in order to establish and maintain communication with the local informants, to identify additional informants, and later to apply quantitative and qualitative tools. The informants from the villages were selected using stratified random sampling. Interviewees in the study all belonged to farmers' communities.

Application of questionnaires, in-depth interviews and semi-structured interviews

Questionnaires were designed to collect information on the main local medicinal plant uses. The questions were answered by a selected group ($n=9$) of old women in Patanoa Rincon in Zetaquirá. Interviews were structured in two groups: in depth interviews and semi-structured interviews, which were done with individual informants in order to maintain data independence [13]. In depth interviews were answered by healers^a and amateur

healers^b. In-depth interviews were carried out in order to collect information on their knowledge on plant use and their services offered to locals. Semi-structured interviews addressed farmers (n = 41). Their distribution across age and sex is shown in Table 1. Semi-structured interviews helped to collect data on traditional plant knowledge and use. For all interviews, vouchers of local medicinal plant species was used as a stimulus for interviewees to elaborate on known and used plants [14].

Homegardens, gardens and solares^c were visited during the interviews. Plant specimens were collected together with the informants when obtainable. From the list of mentioned plants, those that were not cultivated at home were grouped in a separate list.

Disease was defined by interviewees as a period of disequilibrium in the body due to deterioration of organs and functions. Disequilibrium could also be caused by bacteria and viral infection. The diseases described during the study were grouped into categories as it is shown in Table 2.

Field walks

A field walk with key informants was conducted in each municipality in order to identify environmental characteristics of medicinal plant species. The field walk in Campo Hermoso took place along a 5 km transect. In Zetaquirá, it was conducted along a 4.5 km transect. Both transects traversed hills and sloping terrain, agricultural lands, and pastures for cattle ranching with some spots of secondary forests.

Systematics of plant species

Taxonomic identification was mainly conducted by experts at the Colombian National Herbarium (COL) and at the herbarium of Universidad Tecnológica de Tunja and, the Colombian Amazonian Herbarium (COAH) [15]. Collected specimens were deposited at COL, at the non-profit organization Alexander von Humboldt Biological Resources Research Institute Villa de Leyva branch, Colombia (FMB), and at the Herbarium, Botany Group, Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen (CP), for future references. Additionally, specimens were

deposited with the local authorities of each municipality, for reference and further studies at the local level.

Data analysis

Different measures of knowledge described by interviewees can be seen in ethnobotanical research. Therefore, it was decided to carefully evaluate the measure of knowledge to be used in the data analysis of the present study.

Table 2 Categories of diseases with the respective descriptions recorded in the municipalities of Zetaquirá and Campo Hermoso

Category	Diseases
Digestive system	Inflamations of intestine, intestinal obstipation, intestinal obstipation in newborns, diarrhoea, appendicitis, stomach aches, enhance stomach function, gastritis, stomach ulcer, against stomach acids, liver cleanser, hepatitis, against bad breath, againsts umbilical, hernia, diabetes, high bilirubin in new born, tooth pain, mouth driedness, sensitive gums
Bacterial, parasitic and mycotic infections	Inflamations and skin infections, erisipela, againsts acne and feet's fungus, against warts, infections in throat, gripe, intestinal parasites, eye inflammations and infections, eye dust, mouth infections, againsts hair lice, typhus, denge
Musculoskeletal system	Rheumatism, colds and spasms in muscles and tendons, fractures in bones, back pain
Nervous system	Memory loss, nervous system tonics, nervous tension, children with problems to fall asleep, tiredness and weakness, body dryness
Immune system	Fevers, yellow fever
Integumentary system	External and internal inflammations, allergies, infected wounds, excema, spots, rashes in adults, diaper rash
Reproductive system	Prostata complaints, weak uterus, cramps in uterus and menstrual pains, uterus cleanser, menstrual disorders, healing of uterus after giving birth, to prevent extra bleeding during child delivery, to enhance lactation
Circulatory system	Cardiac problems as hypertrophy, hypertension complaints, blood cleanser and tonic, antihemorrhagic, to stop nose bleeding in children
Respiratory system	Sinusitis, to enhance lungs function, asthma, coughs
Urinary system	Children bladder complaints, kidney and bladder complaints in adults
Cancer	Stomach cancer, skin cancer, cancer in general
Accidents	Ofidic accidents
Specific conditions	Headaches, general indisposition, againsts flies, to lose weight, hidropisy, againsts hair loss, hair treatments
Other	Food, condiment herb

Table 1 Distribution of informants who answered semi-structured interviews in the municipalities of Campo Hermoso and Zetaquirá, including totals, percentage of interviewees native to the municipality, and age classes

Municipality	Total	% native	Age class		
			20 to 40	41 to 60	>60
Campo Hermoso	16	81	1	7	8
Zetaquirá	25	80	6	11	8

[illegible]

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics

Species, family, voucher ID, and vernacular names	Life form, Habitat/ place of collection	Status, place of Origin	Index Use Values (estim./ actual)		Part Used	Use	Preparation, mode of administration
			Zeta-quira	Campo Herm.			
<i>Justicia filibracteolata</i> Lindau Acanthaceae ALCG 62 Chuchuhuaza	shrub cultivated in homegarden	native Colombian Andes ³	0.48/0.04	0.06/0.00	bark and leaves (B)	rheumatism (B)	concoction with <i>Allium sativum</i> and <i>Aristolochia ringens</i> in extract in alcohol/ oral -U1- (Z), concoction with <i>Cannabis sativa</i> in extract in alcohol -U2-/ oral (B), decoction/ oral (C)
<i>Trichanthera gigantea</i> (Bonpl.) Nees (s) Acanthaceae ALCG 125 Cafetero, Quiebrabarrigo	small tree wild in edge of route	native to Northern South America ⁸	0.32/0.20	0.19/0.00	leaves and stems (B)	inflammations and skin infections (B) fevers (Z) to lose weight (Z)	decoction/ topical-baths (B) decoction/ topical-baths (Z); fresh extract by maceration/ oral (Z) infusion/ oral (Z)
<i>Sambucus nigra</i> L.* ° Adoxaceae-Caprifoliaceae ++ ALCG 68 Sauco Elder or Elderberry	tree cultivated in solar	introduced Europe-West Siberia-North Africa ⁵	0.72/0.24	0.56/0.25	leaves and flowers (B)* ² flowers (B)° ² leaves and branches (B) flowers and shoots (Z) leaves and branches (B)	gripe (B) eye inflammations and infections (B) fevers (B) tiredness and weakness (Z) respiratory complaints (B) respiratory complaints (Z) headache (Z) against hair lice (C)	concoction with <i>Citrus limon</i> juice in decoction/ oral (B) extract in decoction or destillation (in a glass bottle)/ topical- eye baths (B) decoction/ topical in baths (B) decoction/ topical, baths (Z) decoction/ oral (B) concoction with milk in decoction/ oral (Z) decoction/ oral (B) decoction/ oral (Z) fresh extract from maceration/ topical on the head before the sunrise starts (Z) fresh extract from maceration/ topical for washing the hair (C)
<i>Furcraea macrophylla</i> Baker, Hook (e-CH) Agavaceae Fique,cabuya Fique	succulent rosette forming tall herb cultivated in a farmland	native Boyaca-Colombia ⁷	0.00/0.00	0.25/0.13	root (C) leaves (C)	rheumatism (C) muscles and tendons complaints (C) Hidropisy (C)	decoction/ oral (C) use of fibres of leaves to tie them on the affected part/ topical (C) decoction of a piece of leaf/ oral (C)
<i>Chenopodium ambrosioides</i> L. (s) Amaranthaceae ALCG 80 Paico Epazote Wormseed	herb ascendent cultivated in small garden	native Central Andean range and south of western range in Colombia ⁷	0.40/0.12	0.44/0.38	leaves (B) leaves and stems (B)	stomach complaints (B) intestinal parasites (B)	infusion or decoction/ oral (B) fresh extract by maceration in water/ oral (B)
<i>Althernanthera lanceolata</i> (Benth.) Schinz Amaranthaceae ALCG 113 Planta del cáncer or Sanguinaria	herb cultivated in small garden	introduced American tropics and subtropics, Asia, Africa and Australia ¹²	0.16/0.16	0.48/0.25	leaves and branches (B) leaves (C) leaves (Z)	healing of wounds (B) skin cancer (C) stomach ulcer (C) cancer in general (C) cancer in general (Z) fevers (Z)	decoction/ oral and decoction/ topical-baths (B) decoction/ topical baths (C) decoction in milk/ oral (C) decoction / oral (C) fresh extract by maceration in water/ oral (Z)
<i>Eryngium foetidum</i> L.*	herb cultivated in homegarden	native South Mexico to Bolivia Colombian	0.00/0.00	0.06/0.06	root (C)	hepatitis (C) condiment herb (C)	decoction/ oral (C) combination with food/ oral (C)

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

Apiaceae		Andeans and Orinoquia ⁵			leaves (C)*2		
ALCG 167							
Cilantrón, cilantro cimarrón							
Mexican coriander, shadow beni							
<i>Apium graveolens</i> L. *	herb cultivated in homegarden	introduced Europe and West Asia ¹³ , Europe ⁵	0.20/0.24	0.25/0.19	leaves and stems (B)*2	intestinal obstipation (B)	decoction/oral(B)
Apiaceae							infusion/oral(B)
Apio						stomach ache (B)	combination with food/oral(B)
Celery, celeriac						condiment herb (B)	infusion/oral(B)
						relaxant (B)	decoction/oral(Z)
						to regulate menstruation (Z)	
<i>Foeniculum vulgare</i> Mill.* °	herb cultivated in homegarden	introduced Mediterranean Europe ⁵	0.16/0.16	0.06/0.06	leaves and stems (B)	to enhance lactation (B)	decoction in milk or "aguadepanela"-U3-/oral (B)
Apiaceae							
ALCG 89						condiment herb (B)	combination with food/ oral (B)
Hinojo							
Fennel, Bronze Fennel, Sweet cumin ¹²							
<i>Petroselinum crispum</i> (Mill.) Nyman ex A.W. Hill*	herb cultivated in homegarden	introduced Eastern Mediterranean Western Asia Mediterranean Europe ⁸	0.44/0.36	0.50/0.25	leaves (B)*2	cardiovascular complaints as high blood pressure (B)	infusion/ oral (B)
Apiaceae							to chew leaves/ oral (B)
ALCG 87							combined with food/ oral (B)
Perejil						enhance stomach function (B)	infusion/ oral (C)
Parsley							infusion/ oral (Z)
						againts bad breath (B)	
						condiment herb (B)	
						kidney complaints (C)	
						intestinal pain (C)	
						againts menstrual pain (Z)	
<i>Catharanthus roseus</i> (L.) G. Don.	short lived herb cultivated in pod	introduced Madagascar ⁸	0.00/0.00	0.06/0.06	flowers (C)	againts eye infections and irritations (C)	infusion/ topical-baths (C)
Apocynaceae							
ALCG 158							
Vicaria							
Madagascar periwinkle							

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

<i>Philodendron</i> sp.	sufrutex wild in edges of secondary forest	unknown Africa and Asia and America around the equador ⁷	0.04/0.04	0.00/0.00	leaves and stem (Z)	skin infections and inflamations (Z)	fresh extract by maceration/ topical-rub part affected
Araceae							
ALCG 103							
Hidra or Yedra							
<i>Colocasia esculenta</i> (L.) Schott	herb cultivated in homegarden	introduced South East Asia, Australia ^{8,13}	0.20/0.16	0.31/0.25	tuberous root (B)	gastritis (B)	concoction of blended root with milk/ oral (B)
Araceae						stomach ulcer (B)	
ALCG 106						food (B)	carbohydrate source food/ oral (B)
Malanga							
Taro							
<i>Aristolochia ringens</i> Vahl.	Twiner/climber Wild in river bank	introduced native to Brazil ¹⁵	0.64/0.04	0.75/0.00	leaves and stems (B)	rheumatism (B)	concoction with Cannabis sativa, Erythroxylum coca and Justicia filibracteolata in extract in alcohol/ topical and oral (B), maceration in extract in alcohol/ topical and oral (Z)-U2- maceration/ topical in emplast (B)
Aristolochiaceae						ofidic accidents (B)	
ALCG 78							
Bejuco de Guaco							
Gaping Dutchman`s Pipe							
<i>Aloe vera</i> (L.) Burman f. * °	Short stem-med succulent plant cultivated in pods	introduced arabic peninsula ⁸	0.64/0.60	0.63/0.63	gel from leaves (B) * 2 ° 2	asthma (B)	fresh gel is blended single or in concoction with egg or with honey and juice of Citrus limon/ oral (B)
Asphodelaceae						caugh (B)	
Zábila						fever (B)	
						headache (B)	fresh gel/ topical-emplast over head-front (Z)
						healing wounds and	fresh gel/ topical-emplast over head-front (B)
						skin spots (B)	fresh gel/ topical-emplast over head-front (Z)
Aloe						external inflamations (B)	
						hair treatment (B)	massaging (B)
						stomach cancer (Z)	fresh gel in concoction with honey/ oral (Z)
<i>Bidens pilosa</i> L. *	herb wild in pastures with bushes	native\ From the centrum to the south of the Andeans of Colombia ³	0.00/0.04	0.00/0.00	leaves (Z)*2	skin infections (Z) rashes (Z)	maceration/ topical-emplast (Z)
Asteraceae							decoction/ topical-baths (Z)
ALCG 144							
Chipaca							
Spanish needle,black jack							
<i>Onoseris onoseroides</i> (Kunth) Robins	tall herb wild in slope hills and between bushes on open pastures	introduced Central America and naturalized in Colombia ³	0.00/0.00	0.13/0.00	leaves (B)	against bleedings (C)	fresh leaves directly on the wound/ topical (C)
Asteraceae						against uterus complaints (C)	infusion/ oral (C)
ALCG 139						pains in muscles (Z)	maceration/ topical-emplast (Z)
Santa María							

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

<i>Conyza filaginoides</i> (DC.) Hieron Asteraceae ALCG 110 Venadillo	herb spontaneous in solar with some cultivated trees	native Central Colombian Andean ³	0.04/ 0.00	0.00/ 0.00	leaves and stems (Z)	skin infections and rash (Z)	fresh extract by maceration/ topical-massage (Z)
<i>Pyrethrum parthenium</i> (L.) Sw. * o	herb cultivated in small garden	introduced Eurasian, meridional-boreal ³	0.36/ 0.28	0.00/ 0.00	leaves 2*2 and stems (Z)*2	intestinal complaints (Z) relaxant (Z)	infusion/ oral (Z) fresh maceration/ emplast (Z)
Asteraceae ALCG 75 Manzanilla Feverfew Wild chamomille					flowers (Z)	uterus cramps (Z)	
<i>Galinsoga parviflora</i> Cav. Asteraceae ALCG 90 Guacas	herb cultivated in homegarden	native Colombia, South America ^{3,8}	0.08/ 0.04	0.19/ 0.06	leaves (B) leaves and stems (C)	against excess of stomach acids (B) condiment herb (B) against white plaques in mouth (C) intestinal parasites (C)	decoction/ oral (B) decoction/ oral (B) fresh extract by maceration in water/ oral (C)
<i>Matricaria chamomilla</i> L. * o	herb cultivated in homegarden	introduced Iran and Europe ^{3,8,17}	0.32/ 0.32	0.38/ 0.38	leaves and flowers (B)	stomach complaints (B) relaxant (B) rheumatism (Z)	decoction or infusion (B) infusion/ oral (B) fresh extract by maceration/ topical-massage (Z)
Asteraceae ALCG 96 Manzanilla dulce Roman chamomile, German chamomile					leaves, stems and flowers (B)	colds, spasms and cramps in muscles (C)	decoction/ oral and fresh extract by maceration/ topical-massage (C)
<i>Taraxacum officinale</i> F.H. Wigg. * o	herb spontaneous in a farm with a large patch of secondary forest	introduced Europe ³	0.16/ 0.12	0.25/ 0.06	leaves (B) * 2	cardiovascular complaints (B) kidney complaints (C) enhance lungs function (Z) rheumatism (Z) liver cleanse (Z)	infusion/ oral (B) infusion/ oral (C) infusion/ oral (Z) decoction/ oral (Z) maceration/ topical-emplast (Z)
Asteraceae ALCG 104 Diente de león Dandelion							infusion/ oral (Z)
<i>Artemisia absinthium</i> L. * (e-CH)	herb cultivated in homegarden	introduced Europe ³ and West Asia	0.32/ 0.16	0.19/ 0.13	leaves (B) leaves (Z)	stomach ache (B) liver cleanser (C)	infusion/ oral (B) infusion/ oral (C)

Table 4 List of medicinal plant species reported in Zetaquirá and Campo Hermoso including ethnobotanical characteristics (Continued)

Asteraceae						to prevent extra blood flow during child delivery (Z)	infusion/ oral (Z)
Ajenjo					leaves and stems (Z)		infusion/ oral (Z)
Absinthe wormwood, grand wormwood						against inflammations and infections (Z)	decoction/ topical-baths (Z)
<i>Calendula officinalis</i> L. * °	short-lived plant cultivated in pod	introduced South and East Europe ³	0.60/ 0.24	0.31/ 0.19	flowers (B)	external-internal inflammations (B)	Decoction / topical, baths (B)
Asteraceae					*2 ° 2	gastritis (C)	decoction / oral (B)
ALCG 57					leaves and flowers (C)	skin infections (Z)	infusion of leaves and flowers in concoction with drops of Crotons' sage/ oral (C)
Caléndula							
Pot Marigold, Scotch Marigold					leaves and stems (Z)	menstrual cramps (Z)	maceration of leaves/ topical (Z) decoction/ topical,baths (Z)
<i>Ambrosia cumanensis</i> Kunth	herb cultivated in homegarden	native Colombian Andeans ³	0.88/ 0.40	1.12/ 0.81	leaves and stems (B)	gripe (B)	infusion/ oral (B)
Asteraceae						rheumatism (B)	fresh plant in maceration / topical in cataplasm (B)
ALCG 63					branches (Z)	children bladder complaints (B)	
Altamisa							concoction with <i>Mentha suaveolens</i>
Western Ragweed or Perennial Ragweed						menstrual cramps (Z)	and <i>Ruta graveolens</i> in infusion/ oral (Z)
						colds and spasms in muscles (C)	fresh extract by maceration/ topical to rub body (C)
						stomach ache (C)	decoction/ oral (C)
						intestinal complaints (C)	
						against flies (Z)	branches are placed in dogs beds against flies (Z)
<i>Anredera cordifolia</i> Tenore	Vine wild in edges of secondary forest with enough shadow and moisture	native South America (Colombia, Ecuador, Argentina, Paraguay, Uruguay, Bolivia, Brasil, Salvador) ¹³	0.24/ 0.16	0.12/ 0.13	leaves and stems (B)	fevers (B)	decoction or infusion/ oral (B)
Basellaceae						respiratory complaints (C)	fresh extract by maceration in water/ topical-baths (Z)
ALCG 109							
Rubacá						skin infections (Z)	decoction/ oral (C)
						muscles and tendons inflammations, edemas (Z)	fresh extract by maceration/ topical-emplast or massage (Z)
<i>Crescentia cujete</i> L. ³ ; *	tree cultivated in homegarden	native North Central America, Mexico and Belice, Colombian Andes, Amazonas and Caribbean ³	0.00/ 0.00	0.06/ 0.06	young fruit (C)*2	caught (C)	fresh extract or juice/ oral (C)
Bignoniaceae						asthma (C)	
Totumo							
Calabash tree							
<i>Jacaranda cf. copaia</i> (Aubl.) D.Don	tree cultivated in solar	native Colombian Andeans and Orinoquia ³	0.20/ 0.12	0.94/ 0.69	bark (B)	Intestinal complaints (B)	decoction/ oral (B)
Bignoniaceae					leaves (Z)		infusion/ oral (Z)

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

					leaves (C)	Kidney complaints (Z)	infusion/ oral (C)
ALCG 69					branches (C)	rheumatism (C)	concoction with fresh maceration-Eucalyptus globulus and Allium sativum in extract in alcohol ^{U2} / oral topical (C)
Gualanday						gripe (C)	
						fevers (C)	
						denge (C)	decoction/ oral (C)
						skin infections (C)	see concoction with Urtica dioica and Rosmarinus officinalis (C)
						circulatory complaints (C)	
						mouth dryness (C)	
<i>Symphytum officinale</i> L.	herb cultivated in garden	introduced Europe, West Asia ⁹	0.12/ 0.12	0.06/ 0.13	leaves (B)	rheumatism (C)	infusion/ oral (C)
Boraginaceae						headache (C)	concoction with Rosmarinus officinalis in decoction/ oral (Z)
ALCG 173						prostate complaints (Z)	
Comfrey							
Quaker Comfrey, boneset							
<i>Cannabis sativa</i> L.	herb cultivated	introduced North West of Himalayan mountains, Asia ⁷	0.12/ 0.00	0.19/ 0.06	leaves (B)	rheumatism (B)	concoction with Justicia filibracteolata in extract in alcohol ^{U2} / oral (B),
Cannabaceae							
ALCG 172							concoction with Aristolochia ringens, Erythroxylum coca and Justicia filibracteolata in extract in alcohol/ topical and oral (B)
Marihuana							maceration in extract in alcohol/ topical and oral (Z) ^{U2}
Marijuana,Hemp,Cannabis							fresh fruit juice/ topical directly on skin (C)
<i>Clusia aff. ellipticifolia</i> Cuatr.	small tree cultivated in an edge of pastures plot	native Colombian Andes and Amazons ⁵	0.08/ 0.00	0.13/ 0.00	fruits (C)	against warts (C)	
Clusiaceae or Guttiferae					leaves (Z)	back pain (Z)	heated leaves/ topical directly on the back (Z)
ALCG 122							
Gaque							
<i>Tradescantia zebrina</i> Heynh.	herb cultivated in small garden	introduced Neotropical Guatemala and Mexico ¹³ probably Panamá ⁸	0.04/ 0.08	0.00/ 0.00	leaves and flowers (Z)	diabetes (Z)	decoction/ oral (Z)
Commelinaceae						rheumatism (Z)	maceration/ topical-emplast (Z)
ALCG 115							
Suelda con suelda roja or cinta							
Wandering jew							

Table 4 List of medicinal plant species reported in Zetaquirá and Campo Hermoso including ethnobotanical characteristics (Continued)

<i>Callisia monandra</i> (Sw.) Schult. & Schult. f.	herb wild in edge of secondary forest with enough shadow and moisture	native to Caribbean territories ²⁰ including Colombia	0.12/0.00	0.19/0.13	leaves and stems (B)	fracture bones (B) rheumatism (Z)	maceration/ topical-emplast (B) maceration/ topical-emplast (Z) decoction/ topical-baths (Z)
Commelinaceae							
ALCG 118							
Suelda con suelda blanca							
Cojite morado							
<i>Sechium edule</i> (Jacq.) Sw.	climbing vine plant cultivated in homegarden	introduced Central America and Mexico ⁸	0.08/0.08	0.06/0.00	fruit (B)	hipertension complaints (B)	decoction/ oral (B)
Cucurbitaceae							
ALCG 151							
Guatilla							
Alligator pear,pear squash							
<i>Cupressus lusitanica</i> Mill.	tree cultivated in small garden	introduced Europe ⁷ naturalized in Mexico ²⁴	0.28/0.04	0.13/0.00	branches (B)	rheumatism (B) caughs and asthma (B) against bladder complaints in children (C)	decoction/ topical-baths (B) concoction with Eucalyptus globulus and Solanum nigrum in concentrated extract by decoction/ oral (B)
Cupressaceae							
ALCG 129							
Pino ciprés							
Mexican cypress,Cedar of Goa						tiredness and weakness (Z) kidney complaints (Z)	maceration/ topical-emplast (C) concoction with Eucalyptus globulus in decoction/ topical-baths (Z) decoction/ oral (Z)
<i>Equisetum giganteum</i> L.*	herb wild in strem edge	native Central and South America Central part of Colombian Andeans ⁷	0.28/0.00	0.19/0.00	leaves and stem (B)*2	skin infections and inflamations (B)	decoction/ topical-baths (B)
Equisetaceae							decoction/ oral (Z)
ALCG 156					root (C)	kidney complaints (B)	decoction/ oral (B)
Cola de caballo							
horsetail						againts rheumatism and osteoporosis (C)	concoction with Furcraea macrophylla in decoction/ baths (C)
<i>Erythroxylum coca</i> Lam.	Shrub or small tree Cultivated in solar	native Colombia and South America ⁵	0.08/0.08	0.44/0.25	leaves and branches (B)	rheumatism (B) rheumatism (C)	concoction with Cannabis sativa, Aristolochia ringens and Justicia filibracteolata in extract in alcohol/ topical and oral (B)
Erythroxylaceae							
ALCG 73					leaves (C)	tiredness and	
Coca					leaves (B)	weakness (B)	
					leaves (Z)	relaxant (B) tooth pain (Z)	decoction/ oral (C) infusion/ oral (B) fresh maceration/ topical on the tooth (Z)

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

<i>Croton Funckianus</i> Muell. Arg.	tree wild in edge of route	native Colombian Andes ⁵	0.00/0.00	0.06/0.06	leaves (C)	tooth pain (C)	maceration/ topical-emplast (B)
Euphorbiaceae							
ALCG 120							
Sangregado							
<i>Mimosa pudica</i> L.	herb wild in open pastures	native South and Central America ⁷	0.12/0.12	0.56/0.00	leaves (B)	tooth pain (B)	fresh maceration/ topical-emplast (B)
Fabaceae						rheumatism (C)	decoction/ topical-bath (C)
ALCG 175						for helping children to fall asleep (C)	
Dormidera							
Sensitive Plant, Mimosa							
<i>Senna obtusifolia</i> (L.) Irwin & Barnaby	shrub spontaneous in a sideroad	introduced 48 lower states, Virgin Islands, Puerto Rico ²³	0.12/0.08	0.50/0.06	branches (B)	fever (B)	decoction/ topical-bath (C)
Fabaceae					leaves (C)	gripe (C)	concoction with <i>Trichanthera gigantea</i> and <i>Cestrum mariquitense</i> in decoction/ topical-bath (Z)
ALCG 161					branches (C)	headache (C)	decoction/ oral (C)
Alcaparro						for healing uterus after giving birth (C)	fresh extract by
Sicklepod					branches (B)	to stop noise bleeding in children (Z)	maceration/ topical-massage or bath (C)
							decoction/ topical-bath (C)
							decoction in milk/ topical-bath (Z)
<i>Juglans neotropica</i> Diels (e-CH/Z)	tree cultivated in solar within a secondary forests well drened	native Amazons, Orinoquia Central Andes Range ⁷	0.40/0.20	0.00/0.00	leaves (Z)	allergies (Z)	infusion/ oral (Z)
Juglandaceae					leaves and branches (Z)	diaper rash and other skin complaints as acne and feet's fungus (Z)	decoction/ topical in baths (Z)
ALCG 67							decoction/ topical in hip baths (Z)
Cedro Nogal					leaves (Z)	uterus cleaning (Z)	
Colombian walnut							
<i>Melissa officinalis</i> L.* °	herb cultivated in homegarden	introduced Europe and Asia ³	0.88/0.72	0.75/0.62	leaves*2 and stems (B)*2	gripe (B)	decoction/ oral (B) or fresh extract by
Lamiaceae						fevers (B)	maceration in water/ oral (B) or infusion/ oral (B)
ALCG 92						relaxant (B)	
Toronjil						stomach ache (B)	
Common balm						intestine complaints (B)	
<i>Ocimum campechianum</i> Mill.	herb cultivated in homegarden	native North, Central and Northern South America ²³	0.28/0.16	0.63/0.31	leaves (B)	Intestinal complaints (B)	infusion/ oral (B)

Table 4 List of medicinal plant species reported in Zetaquirá and Campo Hermoso including ethnobotanical characteristics (Continued)

Lamiaceae					seeds (B)	eye dust (B)	one seed is placed in the eye/ topical (B)
ALCG 74						gripe (Z)	infusion/ oral (Z)
Albahaca					leaves (Z)		
Basil, Wild mosquito plant							
<i>Mentha viridis</i> L.	herb cultivated in homegarden	introduced Europe ³	0.36/0.20	0.50/0.38	leaves (B)	gripe (B)	infusion/ oral (B)
Lamiaceae						stomach ache (B)	infusion or decoction/ oral (B)
ALCG 91 Menta						inflammations of intestine (B)	fresh extract by maceration in water/ oral (C)
Spearmint							infusion/ oral (B)
<i>Satureja brownei</i> (Sw.) Briq.	herb cultivated in small garden	native South America, Colombian Andean Ranges ³	0.60/0.20	0.19/0.06	leaves and stems (B)	against bladder complaints in children (B)	maceration/ topical-emplast (B)
Lamiaceae						condiment herb (B)	combination with food/ oral (B)
ALCG 130						stomach ache (Z)	infusion/ oral (Z)
Poleo						enhance blood cleaning (Z)	infusion/ oral (Z)
<i>Origanum majorana</i> L.*	herb cultivated in solar	introduced Asia menor ¹³	0.36/0.28	0.06/0.00	leaves (Z)*2	stomach aches and	infusion or decoction/
Lamiaceae							
ALCG 76						intestine complaints (Z)	oral (Z)
Mejorana						prostate complaints (Z)	infusion/ oral (Z)
knotted marjoram, sweet marjoram						kidney complaints (Z)	infusion/ oral (Z)
<i>Rosmarinus officinalis</i> L.* °	woody herb cultivated in homegarden	introduced Península Ibérica, Western Mediterranean ³	0.44/0.44	0.31/0.19	leaves (B) * 2 ° 2	lungs complaints (B)	infusion/ oral (C)
(e-Z)						tooth pain (B)	decoction in milk/ oral (Z)
Lamiaceae						hipertension complaints (B)	maceration/ topical-emplast on tooth (B)
ALCG 149						against hair loss (B)	infusion/ oral (B)
						muscles pain (C)	fresh extract by maceration/ topical-massage (C)
Romero					leaves and stems (B)	rheumatism (C)	decoction/ topical-baths (Z)
						stomach ache (Z)	fresh extract by maceration/ topical-massage (C)
Rosemary						tiredness (Z)	infusion/ oral (Z)
<i>Mentha suaveolens</i> Ehrh.	herb cultivated in homegarden	introduced Europe ³	1.20/1.04	0.94/0.94	leaves (B)	stomach ache (B)	infusion or decoction/ oral (B)
Lamiaceae					shoots (Z)		fresh extract by maceration/ oral (B)
ALCG 127					leaves and stems (Z)		infusion or fresh extract by maceration/ oral (B)
Yerbabuena blanca					leaves (Z)	headaches (B)	concoction with <i>Allium sativum</i> cloves, in concentrated extract by decoction/ oral (Z)
Round leaved mint					leaves and stems (Z)	intestinal inflammations (B)	

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

						intestinal parasites (Z)	concoction with Piper cf. bogotense, Trichanthera gigantea and Phytolacca rivinoides decoction/ topical-baths (Z)
						against infections and inflammations (Z)	
						fevers (Z)	infusion/ oral (Z)
						menstrual cramps (Z)	concoction with Ambrosia cumanensis and Ruta graveolens in decoction/ oral (Z)
						diaper rash (Z)	decoction/ topical-baths (Z)
<i>Persea americana</i> Mill.	tree cultivated in solar	native Central and South America ⁷	0.04/0.04	0.25/0.13	shoots (C)	gripe (C)	decoction and concoction with fresh extract of Verbena littoralis and Citrus limon juice/ oral (C)
						apendicitis (C)	
Lauraceae					shoots (Z)	sensitive gums (Z)	decoction/ oral (C)
ALCG 100					fruit (C)	against dry hair (C)	maceration/ topical (Z)
Aguacate					fruit (Z)	regulator of digestion (Z)	maceration/ topical-massage (C)
Avocado, butter pear							combination with food/ oral (Z)
<i>Allium fistulosum</i> L.	gregarious herb cultivated in homegarden	introduced Siberia ; Altai montains in Siberia ⁷	0.12/0.08	0.31/0.19	roots (B)	intestinal obstipation (B)	decoction/ oral (B)
							fresh extract by maceration/ oral (C)
Liliaceae					leaves (B)	stomach ache (B)	decoction/ oral (Z)
Cebolla larga					leaves (B)	condiment herb (B)	combination with food (B)
Scallion, green onion					young leaf (C)	against intestinal obstipation in newborns (C)	fresh leaf as suppository/ rectal (C)
					roots (C)	lost of memory (C)	
					leaves (B)	fever (Z)	decoction/ oral (C)
							maceration/ topical-emplast on head-front (Z)
<i>Allium sativum</i> L. * °	herb cultivated in pod	introduced Southwestern Asia; Central Asia ⁷	0.32/0.04	0.12/0.06	bulbus (B) *2 ° 2	gripe (C)	concoction with alcohol-Aguardiente ^{U2} / oral (C)
Liliaceae						fevers (C)	decoction/ oral (C)
Ajo						againsts intestinal worms (C)	concoction with alcohol-Aguardiente ^{U2} / oral (Z)
Garlic						againsts amebiasis (Z)	concoction with shoots of Menta suaveolens in concentrated extract by decoction/ oral (Z)
						rheumatism (Z)	concoction with Justicia filibracteolata and Aristolochia ringens in extract in alcohol ^{U1} - (Z)
<i>Althaea officinalis</i> L. °	herb cultivated in pod	introduced Mediterranean ⁵	0.08/0.04	0.38/0.00	leaves (C)	kidney complaints (C)	infusion/ oral (C)
Malvaceae					flowers (C)	caugh (C)	fresh extract by maceration/ oral (Z)

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

Malva						stomach	infusion/ oral (Z)
Marsh mallow					leaves (B)	ache (Z)	
						fever (Z)	
<i>Anoda cristata</i> (L.) Schl.	herb wild in open pastures	introduced Mexico and part of Central America ⁵	0.00/0.00	0.19/0.13	leaves and stems (C)	fever (C)	concoction with <i>Senna obtusifolia</i> in extract from maceration/ topical-bath or rub body (C)
Malvaceae						healing wounds and againts spots (C)	fresh extract by maceration/ topical-rub skin (C)
ALCG 170							
Patechula (Campo Hermoso)							
Escobo (Zetaquira)							decoction/ topical-bath (C)
Spuded anoda, crested anoda							
<i>Hibiscus rosa-sinensis</i> L.	shrub and groundcover	introduced China ^{11,13}	0.12/0.08	0.31/0.19	leaves (C)	intestinal obstipation (C)	fresh extract by maceration/ oral (C)
Malvaceae	cultivated in small garden				leaves and flowers (C)	fever (C)	concoction with <i>Cestrum mariquitense</i> in decoction/ topical-bath (C)
ALCG 164					leaves (Z)	againts hair loss (C)	decoction/ topical-bath (C)
Rosado					flowers (Z)	nervous system tonic (Z)	fresh extract by blending/ oral (Z)
Hibiscus						eye inflammations (Z)	fresh extract in water/ topical-eye-bath (Z)
<i>Psidium guineense</i> Sw.	small tree spontaneous in a private pastures plot	native Tropical America, native to Colombia ⁵	0.12/0.08	0.44/0.31	leaves and branches (B)	skin infections and recover wounds (B)	decoction/ topical-baths (C)
							maceration/ topical-emplast (Z)
							decoction or infusion/ oral (B)
							infusion/ oral and decoction/ topical-baths (C)
Myrtaceae						intestinal complaints (B)	decoction/ topical-baths (C)
ALCG 121						against umbilical hernia (C)	concoction with other eight bitter plants in decoction/ topical-steam baths (C)
Guayabo Zapatero or Cimarrón						rheumatism (C)	
Brazilian guava, wild guava						against colds and cramps in uterus (C)	
<i>Eucalyptus cf. globulus</i> Labill. * °	tall tree wild in edge of secondary forest	introduced South East of Australia and Tasmania ^{6,10}	0.56/0.00	0.44/0.19	leaves (B)* 2	gripe (B)	infusion/ oral (B)
						fevers (C)	concoction with <i>Jacaranda copaia</i> and <i>Allium sativum</i> in decoction/ topical-baths (C)
						caughs and asthma (B)	concoction with fresh maceration-Jaccaranda copaia and <i>Allium sativum</i> in cloves extract in alcohol ^{U2} / oral (C)

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

						rheumatism (B) tiredness and weakness (Z)	concoction with Solanum nigrum and Cupresus lusitanica in concentrated extract by decoction/ oral (Z) decoction/ topical-baths (B) concoction with Cupresus lusitanica in decoction/ topical-baths (Z)
					° 2 branches (C) leaves (B) branches (B) leaves (Z)		
Myrtaceae							
ALCG 128							
Eucalipto							
Gum tree, Silver-leaved							
mountain gum							
<i>Cattleya schroederiae</i> Rchb. F.	Epiphytic herb cultivated in solar	native Colombia ⁸	0.00/ 0.00	0.69/ 0.63	leaves (C)	typhus (C) denge (C) fever (C) intestinal complaints (C)	fresh extract by maceration of one leafe/ oral (C) extract by blending in concoction with water/ oral (C) fresh extract by maceration of a piece of leafe in concoction with water/ oral (C)
Orchidaceae							
ALCG 165							
Lirio							
Easter orchid, Baron schroeder's							
cattleya							
<i>Petiveria alliacea</i> L. *	herb cultivated in small garden	native Andean (specially Peruvian Amazonas), Caribbean and South east of Colombia ⁷	0.00/ 0.00	0.25/ 0.25	leaves (C)*2	stomach cancer (C) against lacerations in mouth (C) headache (C)	infusion/ oral (C) decoction/ oral (C) extract by maceration/ topical-rinsing mouth out/ (C) infusion/ oral (C)
Phytolaccaceae							
ALCG 136							
Anamú							
Anamú plant							

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

<i>Phytolacca rivinoides</i> Kunth & C.D.Bouché	shrub spontaneous in a solar	native Central and South America, Colombian Andeans ⁷	0.08/ 0.00	0.38/ 0.25	leaves and branches (B)	rheumatism (C) erisipela (C) inflammations and skin infections (C) inflammations and skin infections (Z)	decoction/ topical-baths (C) concoction with <i>Mentha suaveolens</i> , <i>Piper bogotense</i> and <i>Trichanthera gigantea</i> in decoction/ topical-baths (Z)
Phytolaccaceae							
ALCG 146							
Guaba, Cargamanto							
<i>Piper hispidum</i> Sw.	shrub cultivated in solar/it can also occur in wild	native Central and South America and native to Caribbean territories ²⁰	0.00/ 0.00	0.44/ 0.38	leaves and stems (C)	high blood pressure (C) fevers (C) tiredness and weakness (C) rheumatism (C)	fresh extract by maceration in water/ oral (C) concoction with <i>Verbena littoralis</i> in decoction/ topical-bath (C) decoction / topical-bath (C), steam baths/ topical (C)
Piperaceae							
ALCG 85							
Cordoncillo negro							
Jamaican pepper							
<i>Piper cf. bogotense</i> C.DC.	shrub cultivated in solar/it can also occur in wild	native North of South America ¹²	0.08/ 0.00	0.50/ 0.25	leaves (C) leaves and stems (Z)	high blood pressure (C) fevers (C) mouth dryness (C) skin infections and inflammations (Z)	fresh extract by maceration in water/ oral (C) extract in water/ oral (C) concoction with <i>Mentha suaveolens</i> and <i>Phytolacca rivinoides</i> in decoction/ topical- bath (Z)
Piperaceae							
ALCG 81							
Cordoncillo blanco							
<i>Plantago major</i> L.*	herb spontaneous in homegarden	introduced Europe and Asia ³	0.32/ 0.16	0.81/ 0.31	leaves (B)*2	eye dust and infections (B) gastritis (B) liver cleanser (B) kidney complaints (C) fever (C) eczema (C)	decoction/ topical eye bath (C) fresh extract by distillation in glass bottle/ topical-eye drops (Z) fresh extract by maceration/ oral (B) decoction/ oral (C) fresh extract by maceration/ oral (Z) decoction/ oral (C) decoction/ oral (C) fresh maceration in concoction with honey/ topical-emplast (C) concoction with <i>Calendula officinalis</i> and <i>Solanum nigrum</i> in decoction/ topical-baths (Z)
Plantaginaceae							

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

ALCG 374						healing wounds (Z)	
Llantén							
Common plantain							
<i>Cymbopogon citratus</i> (DC.) Stapf.*	herb cultivated in solar	introduced India ²	0.56/0.60	0.44/0.50	leaves (B)*2	stomach aches (B)	infusion or decoction/ oral (B)
Poaceae							infusion/ oral (Z)
							decoction/ oral (Z)
						intestine complaints (Z)	decoction/ oral (C)
						gripe (Z)	
						prostate complaints (C)	
ALCG 77							
Limonaria or limoncillo							
West Indian, Lemon grass, oil grass						fevers (C)	
<i>Rumex crispus</i> L.	herb wild in open pastures	introduced Europe ⁷ , West Asia ²⁰	0.08/0.04	0.19/0.00	leaves (B)	fevers (B)	fresh extract by maceration/ oral (C)
							fresh extract by maceration/ topical-baths (Z)
						kidney complaints (C)	fresh extract by maceration/ oral (C)
						liver cleanser (C)	heated leaves/ topical directly on the lower belly (Z)
Polygonaceae						intestinal inflammations (Z)	
ALCG 148							
Romaza, Lengua de vaca							
Curled dock							
<i>Rubus glaucus</i> Benth.	sub-shrub cultivated in homegarden	native from Mexico to Ecuador and from center to south of Colombian Andeans ⁷	0.28/0.00	0.50/0.13	shoots and fruits (B)	rheumatism (B)	decoction/ oral and decoction/ topical-baths (B)
						against skin infections and inflammations (C)	decoction/ topical-baths and infusion/ oral (C)
							decoction/ topical-baths (C)
Rosaceae					shoots (Z)	against rashes (C)	decoction/ oral (Z)
						against spots (C)	
						caughs (Z)	
ALCG 126							
Moras							
Andean blackberry							
<i>Citrus aurantium</i> var. Amara L.	small tree cultivated in homegarden	introduced Southern Vietnam ¹³ , Southeast Asia ⁵	0.20/0.20	0.69/0.25	fruit (B)	relaxant (B)	steam baths/ topical (B)
							infusion/ oral (C)
							decoction/ topical-for rinsing mouth (B)
						headache (C)	steam baths/ topical (C)

Table 4 List of medicinal plant species reported in Zetaquirá and Campo Hermoso including ethnobotanical characteristics (Continued)

								decoction/ topical-baths (Z)
							mouth infections (B)	decoction/ oral (Z)
							rheumatism (B)	
Rutaceae					leaves (Z)		body dryness (Z)	
Naranjo agrio								
Bitter orange, Seville orange								
<i>Citrus maxima</i> (Burm. ex Rumph.) Merr.	small tree cultivated in homegarden	introduced South East Asia ^{5,13}	0.36/0.24	0.25/0.19	fruits (B)		gripe (B)	juice, alone or in combination with aguadepanela-U3-/ oral (B)
							stomach ache (B)	decoction of shoots/ oral (C)
								juice in combination with water / oral (B)
							inflammations because infections (C)	decoction / topical-baths (C)
								juice in combination with water/ oral (Z)
					shoots (C)		headache (Z)	decoction/ topical-baths (Z)
							fevers (Z)	
					fruits (B)		rheumatism (Z)	
					fruits and branches (C)			
Rutaceae					fruits (Z)			
ALCG 98					fruits and branches (Z)			
Limón mandarin								
Pommelo, Pumelo								
<i>Ruta graveolens</i> L.*	herb cultivated in small garden	introduced Canarian Islands Europe meridional ⁵	1.24/1.08	0.75/0.50	leaves (B)		to strengthen uterus (B)	fresh chopped leaves combined with boiled egg/ oral (B)
								decoction/ oral (B)
								decoction/ topical-bath (Z)
								fresh extract by maceration in water/ oral (C)
Rutaceae					leaves, flowers and stems (B)*2		against cramps in uterus and menstrual pains (B)	infusion/ oral (Z)
							rheumatism (Z)	
							intestinal parasites (C)	
					leaves and stems (C)		cardioregulator (Z)	
ALCG 86					leaves (Z)			
Ruda								
<i>Citrus limon</i> (L.) Burm. F. (e-Z)	small tree cultivated in homegarden	introduced Himalaya and Indochina ⁵	0.52/0.20	1.00/0.63	fruits (B)		gripe (B)	juice alone or in combination with water or aguadepanela/ oral (B)

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

						fevers (B)	juice in combination with water/ oral (B)
						against infections in throat (B)	juice/ to gargle-topical (B)
							juice in combination with water/ oral (B)
Rutaceae						stomach complaints (B)	a fresh hitted-hot fruit/ to massage-topical (C)
							fruit pieces mixed with salt in water/ to do baths-topical (C)
ALCG 93						intestine complaints (B)	fresh fruit pulp/ to rub the head-topical (C)
Limón ácido						rheumatism (C)	juice in combination with water/ oral (Z)
						againts excema (C)	
						headache (C)	
Lemon,Citrus						to prevent high blood presure (Z)	
<i>Physalis peruviana</i> L.	shrub cultivated in homegarden	introduced Northern South America-Peru ³	0.16/ 0.04	0.19/ 0.00	fruits (B)	eye dust and infections (B)	fresh extract from maceration/ topical-eye drops (B)
Solanaceae							a variation is a concoction with honey (C)
ALCG 163							
Uchuba,Guchuba							
Cape gooseberry							
<i>Solanum nigrum</i> L. *	sub shrub spontaneous in solar	introduced Eurasia ^{13,23}	0.20/ 0.04	0.19/ 0.00	fruits and leaves (B)*2	diabetes (C)	infusion/ oral (C)
							decoction/ topical-baths (Z)
						inflammations and skin infections (Z)	concentrated extract by decoction/ oral (Z)
Solanaceae						respiratory complaints (Z)	
ALCG 157							
Yerbamora(Zetaquira)							
Almoraduz(Campo Hermoso)							
black night shade							
<i>Cestrum mariquitense</i> Kunth	shrub cultivated in solar	native Tropical America ¹³ Boyacá and Cundinamarca-Colombia ³	0.08/ 0.08	0.50/ 0.25	leaves (B)	sinucitis (B)	fresh extract by maceration/ topical-noise drops (B)
						fevers (B)	fresh extract by maceration/ oral (B)
						headache (C)	concoction with Hibiscus rosa sinensis in decoction/ topical-baths (C)
Solanaceae							fresh extract by maceration/ oral (C)

Table 4 List of medicinal plant species reported in Zetaquirá and Campo Hermoso including ethnobotanical characteristics (Continued)

ALCG 102							
Tinto							
<i>Cecropia</i> sp.	tree wild in secondary forest	native Colombian Andes ⁷	0.08/0.00	0.06/0.00	leaves (Z)*2	cardiac complaints as hypertrophy (B)	concoction with alcohol-dye/ oral (B)
Urticaceae •							
Cecropiaceae ++							
Yarumo							
Pumpwood, trumpet tree							
<i>Urera baccifera</i> (L.)	herb spontaneous in a farm with a large patch of secondary forest	native Central and South America Central Andeans and Orinoquia in Colombia ⁷	0.20/0.08	0.25/0.06	leaves and branches (B)	rheumatism (B) allergies (B)	decoction/ topical baths (B) infusion/ oral (C)
Gaudich ex Wedd. Urticaceae					leaves (C)	blood circulation tonic (C)	
ALCG 99							
Ortiga mayor, Pringamoza							
<i>Urtica dioica</i> L.* °	herb spontaneous in a farm with a large patch of secondary forest	introduced eurasian, meridional-boreal, native of Britain ¹⁸	0.44/0.32	1.12/0.63	leaves and stems (B)	rheumatism (B) allergies (B)	fresh-maceration/ topical in cataplasm (B) decoction / topical in baths (B) Infusion/ oral (B)
						blood cleaner and tonic (B)	fresh extract by maceration/ topical to rub skin (C) infusion/ oral (C)
Urticaceae					leaves (B)*2	antihemoragic (B) skin infection and inflammations (C)	concoction with Jacaranda copaia and Rosmarinus officinalis in decoction/ oral (C)
					leaves and stems (C)	gripe (C)	
ALCG 66					leaves (C)	mouth dryness (C)	
Ortiga Blanca or Ortiga menor							
Great Nettle or Stinging Nettle							
<i>Lippia alba</i> (Mill.) N.E. Brown*	shrub cultivated in homegarden	native to Colombia ^{5,25}	0.68/0.56	0.75/0.56	leaves (B)*2	stomach complaints (B), diarrhoea (B) tiredness and weakness (C), (Z)	decoction / oral (B) decoction/oral (C),infusion/oral (Z) infusion/oral (Z)
Verbenaceae						childbirth (Z)	
ALCG 72							
Pronto alivio							
Bushy,Matgrass							

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

<i>Lippia citriodora</i> (Lam.) Kunth*	shrub cultivated in homegarden	native Colombian Andean range ⁸	0.48/0.36	0.25/0.13	leaves (B)*2	stomach and intestinal complaints (B)	infusion/ oral (B)
						relaxant (B)	decoction/ oral (B)
						general indisposicion (Z)	infusion/ oral (Z)
Verbenaceae							
ALCG 70							
Cidrón							
Lemon Scented Verbena							
<i>Lantana camara</i> L.	herb wild in open pastures	native to Caribbean territories including Colombia, Continental US and Hawai ²⁰	0.20/0.08	0.25/0.25	leaves and stems (B)	high bilirubin in newborns (B)	decoction/ topical-baths (B)
							infusion/ oral (C)
							decoction in milk/ oral and decoction/ topical-baths (C)
						hepatitis (C)	decoction/ topical-baths (Z)
						yellow fever (C)	
Verbenaceae					flowers (C)	hepatitis (Z)	
					flowers, stems and leaves (C)		
ALCG 117					leaves and stems (Z)		
Florota							
Cambara de Espinto or Pricky Lantana							
<i>Verbena littoralis</i> Kunth *	herb cultivated in solar	native western and central Andean ranges ⁵	0.96/0.40	0.88/0.63	leaves and stems (B)	fevers (B)	decoction/ topical baths (B)
							fresh extract by maceration in water/ oral (Z)
						gripe (B)	concoction with Calendula officinalis and Piper hispidum in decoction/ topical baths (C)
							concoction with Piper hispidum in decoction/ topical in baths (C)
						rheumatism (Z)	infusion/ oral (Z)
						rheumatism (C)	
Verbenaceae						tiredness and weakness (C)	
ALCG 79							
Verbena blanca							
Vervain, owi							
<i>Viola odorata</i> L.*	Acaule herb Cultivated in small garden	introduced Europe, Austral Asia and Boreal Africa ⁵	0.04/0.00	0.00/0.00	leaves*2 and stems (Z)	cough (Z)	infusion/oral (Z)
Violaceae						asthma (Z)	

Table 4 List of medicinal plant species reported in Zetaquira and Campo Hermoso including ethnobotanical characteristics (Continued)

ALCG 111

Violeta

Sweet violet/common
violet

B: Both municipalities; C: Campo Hermoso; Z: Zetaquira.

^{U1} A glass bottle is filled with plant(s) parts, red grape wine usually and, it is buried under the ground during a month.

^{U2} A similar process as it is described in U1, with the variation of the use of aguardiente, which is the introduced alcoholic drink similar to spirit (snaps).

^{U3} Aguadepanela is a traditional colombian homemade drink. The ingredients are water and panela, that is an unrefined food product of which the main component is sugarcane juice.

* Reported in Colombian Vademecum, 2008; * ² Plant material of interest reported in Colombian Vademecum, 2008.

° Reported in WHO, 2009; ° ² Plant material of interest reported in WHO, 2009.

• www.tropicos.org.

++ www.ipni.org/ipni/plantnamesearchpage.do.

(e-CH/Z) Endangered plant species according to the perceptions of the locals in Campo Hermoso (CH) and/or Zetaquira (Z).

¹ Rothmaler Werner, 1994.

² Antolinez González J. C. et.al ,2008.

³ Garcia Barriga Tomo III 1992.

⁴ De Fraume Melida, 1988.

⁵ Garcia Barriga Tomo II, 1992.

⁶ Fonnegra, 2007.

⁷ Garcia Barriga Tomo I 1992.

⁸ Perez-Arbelaes 1978.

⁹ Centro de Educacion No Formal FUNIBA.

¹⁰ Carlos Paez Perez, 1964.

¹¹ www.Botanical.com.

¹² www.zipcodezoo.com/plants/.

¹³ www.en.wikipedia.org.

¹⁴ www.mansfeld.ipk-gatersleben.de.

¹⁵ www.flowersofindia.net.

¹⁶ www.mobot.org.

¹⁷ Soloiki M et al., 2008.

¹⁸ www.botanicus.org (Missouri Botanical Garden).

¹⁹ www.issg.org (global invasive species database).

²⁰ www.cbif.gc.ca/itis (Integr. Taxonomi Inf. Sys.).

²¹ Missouri Bot. Garden 2007 in Colomb. Vademecum.

²² a.o. (tropicalforages.info).

²³ http://plants.usda.gov/java/profile?symbol=SEOB4#.

²⁴ http://www.conifers.org/cu/cup/lusitanica.htm.

²⁵ Vera et. al. in Revista Cubana de plantas medicinales 2010.

²⁶ http://zipcodezoo.com/Key/Plantae/Cecropia_Genus.asp.

²⁷ http://plants.jstor.org/flora/flos003208.

Table 5 Plant species reported as disappeared or endangered within the municipalities of Campo Hermoso and Zetaquirá

	Campo Hermoso	Zetaquirá
Disappeared species	<i>Brownea ariza</i>	<i>Fraxinus udhei</i>
	<i>Ananas comosus</i>	<i>Schizolobium parahybum</i>
	<i>Parietaria officinalis</i>	<i>Chrysophyllum colombianum</i>
	<i>Furcraea</i> sp.	'Alma negra'
	<i>Malva</i> sp.	
	'Gualola'	
	'Mano de León'	
	'Bejuco de roca'	
Endangered Species	<i>Furcraea macrophylla</i>	<i>Citrus limon</i>
	<i>Artemisia absinthium</i>	<i>Saccharum officinarum</i>
	<i>Ocimum campechianum</i>	<i>Juglans neotropica</i>
	<i>Juglans neotropica</i>	<i>Rosmarinus officinalis</i>
	<i>Ceroxylum quindiuense</i>	
	<i>Cedrela</i> spp.	

Wilcoxon-Mann-Whitney tests [16,17] were used to test for differences between introduced and native medicinal plant species in their estimated and actual UV within the two municipalities.

Finally, similarities of use of introduced and native medicinal plant species between the municipalities were calculated using the Jaccard Index [21,22].


Results

Medicinal plant species and plant characteristics


Interviews and questionnaires supplied a list of a total of 80 medicinal plant species used in the treatments of ailments within the municipalities of Campo Hermoso and Zetaquirá. Of these, 78 species were taxonomically identified. Results are based on the 78 identified species of which 35 were native species and 43 introduced (Table 4), belonging to 74 genera and 41 floristic families. The families with the highest numbers of species reported as medicinal were: Asteraceae (10 species), Lamiaceae (7), Apiaceae, Rutaceae and Verbenaceae (each 4), and Malvaceae, Solanaceae and Urticaceae (each 3).


Twenty-nine of the identified species and the genus *Cecropia* Loebl. were included in the list of medicinal plants evaluated and accepted in the Colombian pharmacopeia [23]. Thirteen species were reported in the World Health Organization's (WHO) monographs on selected medicinal plants [24] (Table 4).

During interviews and meetings interviewees were able to reflect about the local resources of medicinal plant species, their importance and, additionally, to discuss




factors that may have increased or reduced plant diversity, such as logging, over- exploitation and difficulties of cultivation. Some interviewees reported that they found it difficult to treat certain diseases because they failed to find the plant species needed. An example is *Brownea ariza* Benth., a native species, which locals consider





extinct in the Campo Hermoso area. This species was used as a haemostat, i.e. to stop bleeding, and as laxative. Another example is *Juglans neotropica* Diels., a native species, used as fungicide and bactericide, which was reported to be endangered in both municipalities (Table 4 and 5). Due to the lack of samples it was not possible to scientifically identify all the species reported as threatened and disappearing by the interviewees.



According to informants' reports and the subsequent calculations of average UVs the most popular medicinal species, in the municipality of Campo Hermoso were *Urtica dioica* (0.88), *Jaccaranda copaia* (0.81) and *Citrus limon* (0.81) while in Zetaquira the most popular species were *Ruta graveolens* (average UV = 1.16), *Melissa officinalis* (0.8), and *Aloe vera* (0.6). Across both municipalities, the most common and popular medicinal plant species were *Mentha suaveolens*, *Ambrosia cumanensis* Kunth, and *Verbena littoralis* (Table 4).


In relation to life form and habitat, the collected plant species could be assigned to 10 categories shown in the Figure 2 (Table 4). Of the native species 40% were found in natural habitats whereas 27% of the introduced species were found as naturalised in the wild.

The naturally occurring and naturalised species were distributed widely within families. For example, Asteraceae (2 herbs), Acanthaceae (1 tree), Aristolochiaceae (1 twiner) and Commelinaceae (2 vines) (Table 4).

Traditional plant use in Campo Hermoso and Zetaquira

Traditional plant uses in the municipalities of Campo Hermoso and Zetaquira are represented in Figures 3, 4, 5, 6. The total number of medicinal plants uses described by informants was higher in more remote Campo Hermoso than in Zetaquira.

Common diseases, such as colds, but also a small number of difficult or complicated diseases, such as cancer, were reported to be treated. The majority of plants mentioned by the informants were reported to be effective in curing the diseases they were applied to. Additionally, most of the informants mentioned the importance



of using folk traditions, e.g. prayers, during treatments in order to ensure the effect of the medicinal plant.

The diseases described by interviewees were grouped in 14 categories (Table 2). The five categories of ailments/diseases with the highest numbers of plants reported in both municipalities were associated with: the digestive system, infections, musculoskeletal, the nervous system, and the immune system. The corresponding most popular medicinal plant species are shown in Table 6. Significant dissimilarities between municipalities occurred in treatments of musculoskeletal, immune and urinary systems, and cancer where the total numbers of plants mentioned in more remote Campo Hermoso was almost double the number of Zetaquira. The category with the highest number of medicinal plant species reported from Zetaquira was 'reproductive system' with 11 plants. There were no plant species specifically reported for the treatment of the two parasitic, epidemic, local diseases of Chagas (caused by *Trypanosoma cruzi*) and Malaria (Paludism). Only two species, *Cattleya schroederiae* Rchb.f. and *Jaccaranda cf. copaia*, were reported for the treatment of Dengue, a viral, epidemic mosquito transmitted disease also affecting these populations (Table 4).

Regarding the plant parts used, leaves were the plant part most frequently reported for remedy preparations, in both municipalities. Other commonly used plant parts were, in descending order of total counts in both municipalities: combination of leaves with stems (36), fruits (16), combination of leaves with branches (9), branches (5), roots (4),

and combination of leaves with flowers (3). Other combinations of different plant parts were mentioned only rarely. The use of seeds was not popular in either of the municipalities (Figure 4).

In both municipalities, similar numbers of applications were reported for *Cymbopogon citratus*, *Plantago major*, *Petroselinum crispum* and *Lippia alba* (Mill.) N.E.Br. These species received the highest scores for application of leaves. Most popular species with the highest applications of leaves were: *Cattleya schroederiae* (4) and *Piper cf. bogotense* C.DC. (3), in Campo Hermoso, and *Origanum majorana* L. (3) in Zetaquira (Table 4).

Eighty different ways of preparation of remedies with medicinal plant species were described. They were grouped according to the type of processing: thermic methods, alcoholic extracts or fresh use. Other ways of preparation of the remedies, such as the combination with food and with other plants (concoction) were also mentioned and thus counted. Practices according to beliefs included the effect of moon or sunlight, esoteric ways and specific ways to treat pets. These additional ways of preparing traditional remedies were included in one group named 'other' (Figure 5).

The most popular way of administration remedies was oral followed by the topical in both municipalities (Figure 6). Among the most popular topical ways of administration, 'baths' yielded the highest scores (13 counts) in both municipalities, 26 in Campo Hermoso and 25 in Zetaquira. Massage was the second most popular in Campo Hermoso, while 'emplast' was the second most popular in Zetaquira.

Healers and amateur healers unanimously reported that bitter plants were denominated 'hot plants' ('plantas calientes', in Spanish), while plants with sweet flavour were denominated 'cold plants' ('plantas frias'). These terms suggested the level of caution with which the medicinal plants should be used. Bitter plants should be used in small doses at low frequency because most of them can produce eye or heart problems. Bitter plants should be used orally only in situations caused by musculoskeletal complaints. Further, they recommended to initially use topical treatments and, in the most general sense, to follow a diet when using medicinal plants. Popular plants used for the purposes outlined above in the localities are described in Table 4.

Table 7 shows the plant species with the highest values obtained in aUV according to the sum of the values from both municipalities. In addition, eUV are also included.

Differences and similarities among plant groups and families

UVs of introduced plant species were significantly higher than native species in the more accessible municipality of Zetaquira (Table 8), while there were no significant

Table 6 Most commonly treated categories of diseases with the most popular plant species used for their treatments

Disease category	No. of plants reported in both municipalities	Popular plant species used in treatments
Digestive system	35	<i>Melissa officinalis</i> <i>Cymbopogon citratus</i> <i>Apium graveolens</i>
Infections from bacteria, parasites or fungi	16	<i>Mentha suaveolens</i> <i>Citrus limon</i> <i>Jaccaranda copaia</i>
Muskuloskeletal system	13	<i>Verbena littoralis</i> <i>Urtica dioica</i> <i>Aristolochia ringens</i>
Nervous system	10	<i>Melissa officinalis</i> <i>Matricaria chamomilla</i> <i>Lippia citriodora</i>
Immune system	9	<i>Aloe vera</i> <i>Melissa officinalis</i> <i>Verbena littoralis</i>

Table 7 Plant species according to the highest actual index use values obtained in both municipalities

Scientific name (Voucher number)	aIUV all	aIUV Zq	aIUV CH	eIUV all	eIUV Zq	eIUV CH	Medicinal uses
<i>Mentha suaveolens</i> Ehrh. * (ALCG 127)	1	1.04	0.94	1.10	1.20	0.94	stomach ache(B), headaches(B), intestinal inflammations(B), intestinal parasites(Z), fevers(Z), menstrual cramps(Z), diaper rash(Z)
<i>Ruta graveolens</i> L. *(ALCG 86)	0.85	1.08	0.50	1.05	1.24	0.75	against cramps in uterus and menstrual pains(B), rheumatism(Z), intestinal parasites(C), cardio-regulator(Z)
<i>Melissa officinalis</i> L. * (ALCG 92)	0.68	0.72	0.63	0.83	0.88	0.75	gripe(B),fevers(B),relaxant(B),stomach ache(B), intestine complaints(B)
<i>Aloe vera</i> (L.) Burman f. **	0.6	0.63	0.63	0.64	0.64	0.63	asthma (B), cough(B), fever(B), headache(B), healing wounds and skin spots(B), external inflammations(B), hair treatment (B), stomach cancer(Z)
<i>Ambrosia cumanensis</i> Kunth (ALCG 63)	0.4	0.81	0.98	0.88	0.88	1.13	gripe(B), rheumatism(B), children bladder complaints(B), menstrual cramps(Z), colds and spasms in muscles(C), stomach ache(C), intestinal complaints(C), against flies(Z)
<i>Lippia alba</i> (Mill.) N.E.Br. (ALCG 72)	0.56	0.56	0.71	0.68	0.68	0.75	stomach complaints(B), diarrhoea(B), tiredness and weakness(C)/(Z), childbirth(Z)
<i>Cymbopogon citratus</i> (DC.) Stapf. * (ALCG 77)	0.6	0.50	0.51	0.56	0.56	0.44	stomach aches(B), intestine complaints(Z), gripe(Z), prostate complaints(C), fevers(C)
<i>Verbena littoralis</i> Kunth (ALCG 79)	0.4	0.63	0.93	0.96	0.96	0.88	fevers(B), gripe(B), rheumatism(Z)/ (C), tiredness and weakness(C)
<i>Urtica dioica</i> L. * (ALCG 66)	0.32	0.63	0.71	0.44	0.44	1.13	rheumatism(B), allergies(B), blood cleaner and tonic (B), anti-haemorrhagic(B), skin infection and inflammations(C), gripe(C), mouth dryness(C)
<i>Citrus limon</i> (L.) Burm. F. * (ALCG 93)	0.2	0.63	0.71	0.52	0.52	1.00	gripe(B), fevers(B), infections in throat(B), stomach and intestine complaints(B), rheumatism(C), eczema (C), headache(C), to prevent high blood pressure(Z)
<i>Jacaranda cf. copaia</i> (Aubl.) D.Don (ALCG 69)	0.12	0.69	0.49	0.2	0.2	0.94	Intestinal complaints(B), Kidney complaints(Z), rheumatism (C), gripe(C), fevers(C), dengue(C), skin infections(C), circulatory complaints(C), mouth dryness(C)
<i>Matricaria chamomilla</i> L. *(ALCG 96)	0.32	0.38	0.34	0.32	0.32	0.38	stomach-complaints(B),relaxant(B), rheumatism(Z),colds, spasms and cramps in muscles (C)
<i>Rosmarinus officinalis</i> L. *(ALCG 149)	0.44	0.19	0.39	0.44	0.44	0.31	lungs complaints(B), tooth ache(B), hypertension complaints(B), against hair loss(B), muscles pain (C) rheumatism(C), stomach ache(Z), tiredness(Z)
<i>Petroselinum crispum</i> (Mill. Nyman ex A. W. Hill*(ALCG 87)	0.36	0.25	0.46	0.44	0.44	0.50	cardiovascular complaints as high blood pressure(B), enhance stomach function(B), halitosis(B), condiment herb (B), kidney complaints(C), intestinal pain(C), against menstrual pain(Z)
<i>Lippia citriodora</i> (Lam.) Kunth (ALCG70)	0.36	0.13	0.39	0.48	0.48	0.25	stomach and intestinal complaints(B), relaxant(B), general indisposition(Z)
<i>Mentha viridis</i> L. * (ALCG 91)	0.2	0.38	0.41	0.36	0.36	0.50	gripe(B), stomach ache(B), inflammations of intestine(B)
<i>Sambucus nigra</i> L. *(ALCG 68)	0.24	0.25	0.66	0.72	0.72	0.56	gripe(B), eye inflammations and infections(B), fevers(B), tiredness and weakness(Z), respiratory, complaints(B), head ache(Z), against hair lice(C)
<i>Cattleya schroederiae</i> Rchb.f. (ALCG 165)	0	0.63	0.27	0	0	0.69	typhoid(C), dengue(C), fever(C), intestinal complaints(C)
<i>Calendula officinalis</i> L. * (ALCG 57)	0.24	0.19	0.49	0.6	0.6	0.31	external-internal inflammations(B), gastritis(C), skin infections (Z), menstrual cramps(Z)
<i>Ocimum campechianum</i> Mill. (ALCG 74)	0.16	0.31	0.41	0.28	0.28	0.63	intestinal complaints(B), eye dust (B), gripe(Z)
<i>Chenopodium ambrosioides</i> L. (ALCG 80)	0.12	0.38	0.41	0.4	0.4	0.44	stomach complaints(B), intestinal parasites(B)
<i>Citrus maxima</i> (Burm. ex Rumph.) Merr. *(ALCG 98)	0.24	0.19	0.32	0.36	0.36	0.25	gripe(B), stomach ache(B), inflammations because infections (C), headache(Z), fevers(Z), rheumatism(Z)
<i>Apium graveolens</i> L. * *	0.24	0.19	0.22	0.2	0.2	0.25	intestinal obstipation(B), stomach ache(B), condiment herb (B), relaxant(B), to regulate menstruation(Z)

Table 7 Plant species according to the highest actual index use values obtained in both municipalities (Continued)

<i>Plantago major</i> L*.(ALCG 374)	0.16	0.31	0.51	0.32	0.32	0.81	eye dust and infections(B), gastritis(B), liver cleanser(B), kidney complaints(C), fever(C), eczema(C), healing wounds (Z)
<i>Citrus aurantium</i> var. <i>amara</i> L. **	0.2	0.25	0.39	0.2	0.2	0.69	relaxant(B), headache(C), mouth infections(B), rheumatism (B), body dryness(Z)

Includes also estimated index use values.

*Introduced plant species; **No specimen collected; (C) Campo Hermoso; (Z) Zetaquira; aIUUV Actual Index Use Value; eIUUV Estimated Index Use Value; all-aIUUV/eIUUV Sum of all values corresponding to actual or estimated uses.

differences between the two groups in the more remote municipality of Campo Hermoso and nor in the two municipalities combined.

Numbers of plant species for which estimated and actual UVs could be calculated did not differ greatly between the two municipalities (Table 9). Nevertheless, totals of actual UV of introduced species were higher in Zetaquira than in Campo Hermoso. Totals of actual UV of native species were higher in Campo Hermoso than in Zetaquira. Totals of common plant species including introduced and native ones were very high in both municipalities.

There were many similarities in the use of both introduced and native plant species between the municipalities (Table 10).

Estimated Family Use Values (FUV) were higher than actual family use values in almost all cases (Table 11). Estimated and actual FUV were commonly higher for Campo Hermoso than for Zetaquira. Plant families represented by high numbers of plant species did not always have high FUVs, i.e. Asteraceae and Apiaceae. Some families, of which only one plant species was reported, surprisingly obtained high values, e.g. Aristolochiaceae which had the highest estimated use value. Another interesting case was the Asphodelaceae which yielded high actual use values. Furthermore, Asphodelaceae and Poaceae coincided in having high actual use values in both municipalities.

The list of the most popular medicinal plant species based on median of actual and estimated Use Values, among amateur healers and healers is shown in Table 12.

From the amateur healers' list, 87% species were introduced including six herbs and one shrub, while only 13%, i.e. one herb, was native. In the case of healers, 57% in the list of preferred species were native, which included two trees and two shrubs, while the remaining 43% were introduced, i.e. two herbs and one vine (Table 4).

Discussion

Significance of the use of medicinal plants within the municipalities

The patterns in traditional knowledge and medicinal plant use described in this study is in accordance with other studies [25-27] in the high number of introduced species mostly herbs. Especially for the more accessible municipality of Zetaquira the use of native species was rarely reported.

Rapid adoption of introduced species in traditional medicine can be understood as a response to new opportunities arising with globalisation. Inefficacy of native species may lead to experimentation with introduced species. Introduced species enrich the arsenal of species used in treatments and are often seen as very powerful [28-30]. Adoption of introduced species may be seen as a way to reshape and re-vitalise traditional practices, which in many places provide an important alternative to the official health care services within developing countries.

The effectiveness of treatments with popular introduced plant species that are used world-wide indicates the need to promote and further the studies on the use and effectiveness of these plant species [23,24]. In the Colombian pharmacopoeia list of officially accepted plants for medicinal use 44% of the total of 149 species are introduced [23,31].

The flip-side of the coin is that with the adoption of introduced medicinal plant species peoples' interest in cultivating native species decreases. In this study most of the popular medicinal plant species (UV >0.5) were introduced with the exception of *Verbena littoralis* and *Ambrosia cumanensis* (Table 4 and 7). Furthermore, a number of native species with multiple uses, e.g. timber but also medicinally, are currently endangered or have disappeared from the regions indicating that over-exploitation has taken place [27].

Table 8 Results of Mann-Whitney-Wilcoxon tests of differences in Index Use Values (IUUV) among native and introduced medicinal plants. Med. = Median; Nat. = Native; Int. = Introduced

Index	Campo Hermoso			Zetaquira			Both Municipalities		
	Med. Nat.	Med. Int.	P	Med. Nat.	Med. Int.	P	Med. Nat.	Med. Int.	P
Actual IUUV	0.13	0.13	0.677	0.04	0.14	0.005	0.09	0.12	0.095
Estimated IUUV	0.25	0.25	0.750	0.12	0.20	0.047	0.22	0.23	0.237

Table 9 Numbers of introduced and native medicinal plant species for which estimated and actual Index Use Values could be calculated

Origin	Index Use Value	Campo Hermoso	Zetaquira	Both
Introduced	Actual	4	10	25
	Estimated	2	3	38
Native	Actual	11	5	18
	Estimated	7	2	26

The high similarities between estimated and/or actual UV (> 50%) for introduced and native plant species showed little difference in the lists of medicinal plants between the municipalities. For introduced species, similarity values were the highest (Table 10). These results are comparable to the findings of De Almeida et al. [28] who worked with rural communities in the Northeast Brazil.

Overexploitation of medicinal plant species, including a large number of native species, has been discussed extensively. Njoroge et al. [32] found that *Carissa edulis* (Forssk.) Vahl, a native wild species of Ethiopia [33], is a priority species used in the treatment of several ailments, especially for stomach pains, a common disorder in the Mwingi District, Kenya. Presently, this species is threatened by overuse. Similarly, other scholars have reported that plant species most popularly used by communities depend on the kind of local diseases that people face [2], which could be a parameter to identify possible species that could become endangered by overexploitation in combination with knowledge on plant parts used, harvest techniques, demand and prices.

The most frequent diseases in the study sites according to local reports from the health centre service [34] were related to the digestive system and infections caused by bacteria, parasites or fungi. The present study shows that the highest valued species were related to exactly these diseases. Of these, the top five were all introduced (Tables 4 and Table 7) confirming that communities make use of specific plant species including introduced ones.

The large number of plant species used in the treatment of complaints of the digestive system, infections and nervous system in both municipalities, is comparable to other findings from studies involving farmer communities in other localities in department of Boyacá.

Table 10 Jaccard-index similarity (%) between Campo Hermoso and Zetaquira municipalities for medicinal plant species with estimated and actual Index Use Values

Index Use Value	Introduced	Native
Estimated	88	74
Actual	64	53

For example, Lagos [9] found that stomach-complaints occupy the category of illnesses which is treated with the highest number of medicinal plant species, followed by complaints of the nervous system. At the same time, these results are comparable to findings from other places. In the municipalities of the city of Imbituba, Santa Catarina, Brazil Zank and Hanazaki [35] found, that digestive complaints were the disease category with the highest therapeutic applications of medicinal plants. Furthermore, Neves et al. [36], who studied various groups of the community of Trás-os-Montes, Portugal, including farmers, reported that illnesses related to the digestive system were most popularly treated with medicinal plants.

Life form, habitat and proximity to cities facilitate popularity of introduced species

It has been demonstrated in several studies [28,37,38], that life form and habitat specification determine the use of medicinal plant species. The reason usually mentioned is that herbs, of which a large number are introduced, are easy to cultivate and maintain in small gardens or pots near to or in the houses. Examples of species whose popularity can be related to life form and the ease of cultivation are *Ruta graveolens* that has a high medicinal value in Zetaquira, and *Urtica dioica* that is a highly valuable herb in Campo Hermoso. Opposed to the introduction of easily cultivated herbs, a number of ecological factors diminish advantages of cultivation and maintenance of some native plant species.

Among the most popular species of amateur healers (eight species), seven species (87.5%) were introduced, of which six were herbs and only one a shrub (Table 12). In contrast, among healers the tendency to use native species, usually trees or shrubs, is higher: 57% of their most popular species was native and included two herbs and one vine. Since healers are specialised in the use of medicinal plants, they are in most cases willing to cultivate a diversity of species, including trees and shrubs with medicinal values, in their home gardens and on other cultivated lands. Hence, cultivation could explain the popularity of native woody species within the group of healers.

The high significance of introduced plant species according to UV (Table 8) in Zetaquira closer to the capital of the department, Tunja, could be an indication of the influence of distance to modern cities and acculturation upon the adoption of new medicinal plant species. These findings are comparable to results obtained by other studies, e.g. in Brazil, Argentina and Manus Island (Papua New Guinea) [28,39,40]. Inhabitants of Campo Hermoso, which is more remote and follows rural traditions, have higher knowledge on medicinal plant use. Furthermore, there are more healers in Campo Hermoso. Hence, the null

Table 11 Family Use Values (FUV) of most popular plant families calculated from estimated and actual Index Use Values of species and number of native and introduced species per family

Family	Native	Introd.	FUV est. CH	FUV est. Zq	FUV est. Both	FUV act. CH	FUV act. Zq	FUV act. Both
Asteraceae	4	6	0.26	0.28	0.27	0.16	0.16	0.16
Lamiaceae	2	5	0.48	0.59	0.55	0.36	0.43	0.40
Apiaceae	1	3	0.22	0.20	0.21	0.14	0.19	0.17
Rutaceae	0	4	0.67	0.58	0.62	0.39	0.43	0.41
Verbenaceae	4	0	0.53	0.58	0.56	0.39	0.35	0.37
Malvaceae	0	3	0.29	0.07	0.15	0.10	0.04	0.07
Solanaceae	1	2	0.29	0.15	0.20	0.08	0.05	0.07
Urticaceae	2	1	0.48	0.24	0.33	0.23	0.13	0.17
Acanthaceae	2	0	0.13	0.40	0.30	0.00	0.12	0.08
Alliaceae	0	2	0.22	0.22	0.22	0.13	0.06	0.09
Amaranthaceae	1	1	0.44	0.28	0.34	0.31	0.14	0.21
Bignoniaceae	2	0	0.50	0.10	0.26	0.38	0.06	0.18
Commelinaceae	1	1	0.09	0.08	0.09	0.06	0.04	0.05
Myrtaceae	1	1	0.44	0.34	0.38	0.25	0.04	0.12
Phytolaccaceae	2	0	0.31	0.04	0.15	0.25	0.00	0.10
Piperaceae	2	0	0.47	0.04	0.21	0.31	0.00	0.12
<u>Adoxaceae</u>	0	1	0.56	0.72	0.66	0.25	0.24	0.24
<u>Aristolochiaceae</u>	0	1	0.75	0.64	0.68	0.00	0.04	0.02
<u>Asphodelaceae</u>	0	1	0.63	0.64	0.63	0.63	0.60	0.61
<u>Plantaginaceae</u>	0	1	0.81	0.32	0.51	0.31	0.16	0.22
<u>Poaceae</u>	0	1	0.44	0.56	0.51	0.50	0.60	0.56

Bold = plant families with higher number of species (>2); italics = plant families with two plant species; underlined = plant families with one species but high FUV. Est. = estimated; act. = actual; CH = Campo Hermoso; Zq = Zetaquira; both = both municipalities together.

hypothesis of similarity between plant knowledge and use between municipalities is falsified.

High presence of pharmacologic components in herbs and popularity of genera

The high presence of pharmacologic components in herbs makes them attractive for treatments of different diseases [27]. Specifically the high composition of alkaloids in leaves facilitates medicinal uses as has been indicated by the popularity of these plant parts within

Table 12 Most popular species among healers and amateur healers

Amateur healers	Healers
<i>Calendula officinalis</i> L.	<i>Jacaranda copaia</i> (Aubl.) D.Don
<i>Citrus Limon</i> (L.) Burm.f	<i>Aristolochia ringens</i> Vahl.
<i>Petroselinum crispum</i> (Mill.) Nyman ex A.W. Hill	<i>Ruta graveolens</i> L.
<i>Chenopodium ambrosioides</i> L.	<i>Lippia alba</i> (Mill.) N.E. Brown
<i>Ruta graveolens</i> L.	<i>Althernanthera lanceolata</i> Benth.
<i>Cymbopogon citratus</i> (DC.) Stapf	<i>Psidium guinense</i> Sw.
<i>Mentha suaveolens</i> Ehrh.	<i>Rubus glaucus</i> Benth.
<i>Allium sativum</i> L.	

several communities [9,25,27,36] and also in the present study. Furthermore, a number of phytochemical studies have proven remarkable alkaloid and oil content in herbal leaves [41-43] which may provide alternatives for the pharmaceutical industry. Taxa with large numbers of useful herbs contribute to the tendency of using leaves and introduced plant species. For example, the genus *Mentha*, one of the most popular taxa in Campo Hermoso and Zetaquira and, was reported to be used in treatments of digestive complaints, colds, fevers, skin infections, inflammations and headaches (Table 4). *Mentha ssp.*, introduced to the American continent, is widely popular in folk medicine [9,35,44] and contribute to the popularity of introduced plant species.

Popularity of plant families

Popularity may also reflect world-wide presence of large plant families, such as the Lamiaceae. Contrary, Schippmann et al. [1], suggest that popularity of plant families is related to the local availability of the species. Studies in diverse localities as for example in Loja province, Ecuador [45], Imbituba, Santa Catarina, Brazil [35], Sierras de Córdoba, Argentina [46], and Banbungo, Cameroon [27] have found Asteraceae and Lamiaceae with high

numbers of plant species used locally. These findings are comparatively similar to the results obtained by ethnobotanical studies carried out in the department of Boyacá, e.g. the studies of Lagos [9] in five municipalities in the central part of Boyacá, and a study carried out in the municipality of San José de Pará by Toscano [10]. The results indicate a large availability of species belonging to the Asteraceae and Lamiaceae in this region. At the same time, these results are comparable to the global pattern suggested by Moerman [47] who considered that Asteraceae and Lamiaceae are among the most used plant families in traditional medicine world-wide.

In relation to the popularity of Asteraceae, Garcia Barriga [48] reported that this family is the taxonomic group of plant species with the highest use in traditional medicine in Colombia. Nevertheless, the large range of distribution of this family may explain part of the popularity of this taxonomic group in folk medicine [27].

Furthermore, plant families, such as the Apiaceae, Apocynaceae and Guttiferae, of which there are examples of species with medicinal use in Campo Hermoso and Zetaquirá, are part of the list of plant families with high numbers of traditionally used species worldwide [1].

However, 85% of the medicinal species identified in this study have been described and reported as medicinal in Colombia [11,48-50]. Of these species, 54% are introduced. This indicates that transmission of knowledge on introduced medicinal species has been successful in Colombia and in the studied municipalities. Similar situations have been observed and discussed in other countries as Brazil [28] where it was observed that knowledge of introduced plant species within farmers communities has been well established.

Promotion, conservation and sustainable use of native medicinal plant species may be facilitated by including shrubs and trees into local agroforestry systems. A number of these species, apart from being medicinal, offer also other kind of uses, e.g. as food, source of wood and shade. Such multiuse plants are often valuable to locals [31].

Conclusions

A total of 80 medicinal plant species were recorded. Of these, 78 species were taxonomically identified of which 35 were native species and 43 were introduced belonging to 74 genera and 41 floristic families. Among the native species 40% were found in natural habitats while twenty-seven per cent of introduced species were found as naturalised in the wild. The families with the highest numbers of species reported as medicinal were: Asteraceae, Lamiaceae, Apiaceae, Rutaceae, Verbenaceae, Malvaceae, Solanaceae and Urticaceae. Most applications of medicinal plants are related to diseases of the digestive and respiratory systems, and infections. Leaves are the most popular plant part

used. Decoctions and oral administration are the most common practices.

The total of medicinal plants uses described by informants was higher in the more remote Campo Hermoso than in the more accessible Zetaquirá. UVs of introduced plant species were significantly higher than native species in the more accessible municipality of Zetaquirá, while there were no significant differences between the two groups in the more remote municipality of Campo Hermoso and nor in the two municipalities combined.

The list of the most popular medicinal plant species for healers and amateur healers respectively showed that 87% of the species from the amateur healers' list were introduced including mainly herbs while in the case of healers, 57% of the species of mainly trees and shrubs. Conversion of forests for agriculture and cattle ranching is depleting local forest resources and a number of medicinal plants were reported as disappearing or locally extinct. Only about 50% of the native medicinal species were found in cultivation. Agroforestry may provide an option for integrating agriculture with cultivation of native trees, shrubs and lianas, not otherwise cultivated. Educational programs could raise awareness in relation to conservation and maintaining use of native species.

Consent

Written informed consent was obtained from the patient for publication of this report and any accompanying images.

Endnotes

^a Group of experts on the use of local medicinal plant species who offer local people treatments using medicinal plants.

^b Group of people who are interested in the use of medicinal plants, but do not work as professional healers.

^c Local term to refer to large home gardens with inclusion of a variety of trees and shrubs.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Ana Lucía Cadena-González was the main responsible for study design, data analysis, interpretation, and writing. She conducted the field work in Campo Hermoso and Zetaquirá between October 2008 and February 2009, collected the plant specimens, and identified the specimens in cooperation with the herbaria. Marten Sørensen and Ida Theilade contributed to the concept and design of the study, interpretation of the findings and preparation of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors acknowledge the collaboration of the local communities and authorities from the municipalities of Campo Hermoso and Zetaquirá. We especially thank the interviewees and the participants in community meetings. The staff of the Colombian National Herbarium and of the Herbarium at Universidad Pedagógica y Tecnológica de Colombia is gratefully acknowledged for their help in the identification of the plant

material. Finally, we thank DANIDA and the OTICON fund for the grants supporting the field work.

Author details

¹Department of Agriculture and Ecology, University of Copenhagen, Rolighedsvej 21, 1958 Frederiksberg C, Denmark. ²Department of Plant and Environmental Sciences, University of Copenhagen, Rolighedsvej 21, 1958 Frederiksberg C, Denmark. ³Department of Food and Resource Economics, Faculty of Science, University of Copenhagen, Rolighedsvej 23, 1958 Frederiksberg C, Denmark.

Received: 7 May 2012 Accepted: 28 March 2013

Published: 11 April 2013

References

- Schippmann U, Leaman DJ, Cunningham AB: *Impact of cultivation and gathering of medicinal plants on biodiversity: global trends and issues, Biodiversity and the Ecosystem Approach in Agriculture, Forestry and Fisheries. Satellite event on the occasion of the Ninth Regular Session of the Commission on Genetic Resources for Food and Agriculture: 12–13 October 2002; Rome.* FAO: FAO; 2002:1–21.
- Muthu C, Ayyanar M, Rajan N, Ignacimuthu S: **Medicinal plants used by traditional healers in Kaancheepuram District of Tamil Nadu.** *India. J Ethnobiol Ethnomed* 2006, **2**:43.
- ProFound Advisors in Development: **Market brief in the European Union for selected natural ingredients derived from native species.** *Crescentia cujete L., Calabash, Totumo.* In *Series of Market briefs on selected natural ingredients derived from native species in beneficiary countries of the BTFP. UNCTAD-United Nations Conference on Trade and Development, BioTrade Initiative/BioTrade Facilitation Programme (BTFP).* Edited by United Nations. United Nations; 2005:1–40.
- Dunmire WW: *Gardens of New Spain.* Austin: How Mediterranean plants and foods changed America. University of Texas Press; 2004.
- Quiroga R, Meneses L, Bussmann RW: **Medicinal ethnobotany in Huacareta (Chiquisaca, Bolivia).** *J Ethnobiol Ethnomed* 2012, **8**:29. doi:10.1186/1746-4269-8-29.
- Luziatelli G, Sørensen M, Theilade I, Mølgaard P: **Asháninka medicinal plants: a case study from the native community of Bajo Quimiriki, Junín.** *Peru. J Ethnobiol Ethnomed* 2010, **6**:21.
- Dueñas M: *A: Plan de desarrollo municipal 2008–2011.* Alcaldía Municipal Campo Hermoso-Boyacá: La unidad de compromiso social; 2008.
- Pacheco FP: *Plan de Desarrollo Municipal de Zetaquirá Boyacá 2004–2007.* Desarrollo Municipal con Participación Ciudadana: Alcaldía Municipal Zetaquirá- Boyacá; 2004.
- Lagos LM: **Especies con propiedades medicinales encontradas en las colecciones del herbario UPTC y su uso actual en Tunja y en los municipios de Chiquiza, Cómbita, Oicatá, Sora y Soracá (Boyacá).** *Revista De La Asociación Colombiana De Herbarios. Universidad del Quindío* 2005, **25**:58–68.
- Toscano GJY: **Uso tradicional de plantas medicinales en la vereda San Isidro, municipio de San José de Paré-Boyacá: Un estudio preliminar usando técnicas cuantitativas.** *Acta Biológica Colombiana* 2006, **11**:137–146.
- Perez AE: *Plantas Útiles de Colombia. Tercera redacción.* Bogotá: Litografía Arco; 1956.
- Cunningham AB: *Applied Ethnobotany – People, Wild Plant Use and Conservation.* London: Earthscan Publications; 2002.
- Martin GJ: *Ethnobotany, A methods manual. A People and Plants Conservation Manual.* London: Chapman and Hall; 2004.
- Thomas E, Vandebroek I, Van Damme P: **What Works in the Field? A Comparison of Different Interviewing Methods in Ethnobotany with Special Reference to the Use of Photographs.** *Econ Bot* 2007, **61**:376–384.
- Colombian Amazonian Herbarium (COAH). www.sinchi.org.co/herbario.
- Dytham C: *Choosing and Using Statistics – A Biologist's Guide.* Oxford: Blackwell Sciences; 1999.
- Quinn GP, Keough MJ: *Experimental Design and Data Analysis for Biologists.* Cambridge: Cambridge University Press; 2002.
- Sánchez M, Duivenvoorden JF, Duque A, Miraña P, Cavellier J: **A Stem-based Ethnobotanical Quantification of Potential Rain Forest Use by Mirañas in north west Amazonia.** *Ethnobot Res Appl* 2005, **3**:215–229.
- Phillips O, Gentry A: **The Useful Plants of Tambopata, Peru: I. Statistical Hypotheses Tests With a New Quantitative Technique.** *Econ Bot* 1993, **47**:15–32.
- Hoffman B, Gallaher T: **Importance Indices in Ethnobotany.** *Ethnobot Res Appl* 2007, **5**:201–218.
- Mc Cormick WP, Lyons NI, Hutcheson K: **Distributional properties of Jaccard's Index of similarity.** *Commun Stat-Theor M* 1992, **21**:51–68.
- Campos MT, Ehringhaus C: **Plant Virtues are in the eyes of the beholders: A comparison of Known Palm Uses among Indigenous and Folk Communities of South-Western Amazonian.** *Econ Bot* 2003, **57**:324–344.
- Ministerio de Protección Social, Colombia: *Vademecum colombiano de plantas medicinales.* Bogotá, D.C: El Ministerio; 2008.
- World Health Organization: *Monographs on selected medicinal plants.* Geneva, Switzerland: WHO Press; 2009.
- Estupiñán GA, Jiménez EN: **Uso de las plantas por grupos campesinos en la franja tropical del Parque Nacional Natural Paramillo (Córdoba, Colombia).** *Caldasia* 2010, **32**:21–32.
- Ladio AH, Lozada M, Weigandt M: **Comparison of traditional wild plant knowledge between aboriginal communities inhabiting arid and forest environments in Patagonia, Argentina.** *J Arid Environ* 2007, **69**:695–715.
- Simbo DJ: **An Ethnobotanical Survey of medicinal plants in Babungo, Northwest Region.** *Cameroon. J Ethnobiol Ethnomed* 2010, **6**:8.
- De Almeida CC, Branco R, Ramos AM, De Amorin C, De Albuquerque UP: **A comparison of knowledge about medicinal plants for three rural communities in the semi-arid region of north-east of Brazil.** *J Ethnopharmacol* 2010, **127**:274–284.
- De Albuquerque UP: **Re-examining hypotheses concerning the use and knowledge of medicinal plants: a study in the Caatinga vegetation of NE Brazil.** *J Ethnobiol Ethnomed* 2006, **2**:30.
- Theilade I, Hansen HH, Krog MP: **Ethnobotanical knowledge: implications for participatory forest management.** *J Transdiscipl Environ Stud* 2007, **6**:1–14.
- Fonnegra GR: **Antecedentes Para la Construcción de una Farmacopea ó el Vademecum Colombiano de Plantas Medicinales.** In *Corporchivor, Instituto Humboldt, Universidad Pedagógica y Tecnológica de Colombia.* Universidad Pedagógica y Tecnológica de Colombia: Universidad Distrital Francisco José de Caldas; 2006:101–160.
- Njoroge GN, Joroge GN, Kaibui IM, Njenga PK, Odhiambo PO: **Utilisation of priority traditional medicinal plants and local people's knowledge on their conservation status in arid lands of Kenya (Mwingi District).** *J Ethnobiol Ethnomed* 2010, **6**:22.
- CRAF-PROSEA network: *World Agroforestry Centre, Agroforestry Tree Database.* http://www.worldagroforestry.org/treedb2/AFTPDFS/Carissa_edulis.pdf.
- Secretaría de Salud de Boyacá: *Análisis Situacional del Municipio de Zetaquirá.* Boyacá: Dirección Técnica de Salud Pública, Vigilancia en Salud Pública; 2008.
- Zank S, Hanazaki N: *Exploring the links between Ethnobotany, Local Therapeutic Practices, and Protected Areas in Santa Catarina Coastline.* Evid Based Complement Alternat Med: Brazil; 2011. doi:10.1155/2012/563570.
- Neves JM, Matos C, Mouthinho C, Queiroz G, Gomes Lígia Rebelo G: **thnopharmacological notes about ancient uses of medicinal plants in Trás-os-Montes (northern of Portugal).** *J Ethnopharmacol* 2009, **124**:270–283.
- De Albuquerque UP, Da Silva VA, Cabral M, Alencar NL, Andrade L: **Comparisons between the use of medicinal plants in indigenous and rural caatinga (dryland) communities in NE Brazil.** *Press Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromaticas* 2008, **7**:156–170.
- Merlin F, Narasimham D: **Plant names and uses as indicators of knowledge patterns.** *Indian J Tradit Know* 2009, **8**:645–648.
- Ladio AH, Lozada M: **Comparison of wild edible plant diversity and foraging strategies in two aboriginal communities of northwestern Patagonia.** *Biodivers Conserv* 2003, **12**:937–951.
- Case RJ, Pauli GF, Soejarto DD: **Factors in maintaining indigenous knowledge among ethnic communities of Manus Island.** *Econ Bot* 2005, **59**:356–365.
- Johnson E, Emche S: **Variation of Alkaloid Content in *Erythroxylum coca* Leaves from Leaf Bud to Leaf Drop.** *Ann Bot-London* 1994, **73**:645–650.
- Barni M, Fontanals A, Moreno S: **Estudio de la eficacia antibiótica de un extracto etanólico de *Rosmarinus officinalis* L. contra *Staphylococcus aureus* en dos modelos de infección en piel de ratón. [Study of the antibiotic efficacy of an ethanolic extract from *Rosmarinus officinalis***

against *Staphylococcus aureus* in two skin infection models in mice]. *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas* 2009, **8**:219.

43. Rojas LB, Usubillaga A: **Composition of the essential oil of *Satureja brownei* (S.W.) Briq. from Venezuela.** *Flavour Frag J* 2000, **15**:21–22.
44. Gomez H, Díaz F, Franco L, Mercado J, Guzmán J, Medina J, Gaitán R: **Folk medicine in the northern coast of Colombia: an overview.** *J Ethnobiol Ethnomed* 2011, **7**:27.
45. Morochó V, Tinitana F, Malagón O, Zaragoza T, Armijos C, Medina M, Medina I, Chalán D, Lojan C, Lavanda C, Minga P, Cartuche J: **Estudio Etnobotánico de especies medicinales en la comunidad indígena "Saraguro" de la Provincia de Loja, Ecuador.** In *Proceedings of III National Conference in Medicinal and Aromatic Plants, and Tropical Floriculture: 26–27 October 2006; Municipality of Jenesano, Boyacá*. Edited by Corpochivor: Instituto Humboldt. Universidad Distrital Francisco José de Caldas: Universidad Pedagógica y Tecnológica de Colombia; 2006:46–80.
46. Martínez GJ, Luján MC: **Medicinal plants used for traditional Veterinary in the Sierras de Córdoba-Argentina: An ethnobotanical comparison with human medicinal uses.** *J Ethnobiol Ethnomed* 2011, **7**:23.
47. Moerman DE, Pemberton RW, Kiefer D, Berlin B: **A comparative analysis of five medicinal floras.** *Journal of Ethnobiol* 1999, **19**:49–67.
48. García Barriga H: *Flora Medicinal de Colombia, Tomos I–III*. 2nd edition. Bogotá: Tercer Mundo Editores; 1992.
49. Pbro. Arias E: *Plantas Medicinales*. Vigésima Ediciónth edition. Medellín: La Pluma de Oro; 1991.
50. Fonnegra R, Jimenez SL: *Plantas medicinales aprobadas en Colombia*. 2nd edition. Medellín: Universidad de Antioquia; 2007.

doi:10.1186/1746-4269-9-23

Cite this article as: Cadena-González et al.: Use and valuation of native and introduced medicinal plant species in Campo Hermoso and Zetaquirá, Boyacá, Colombia. *Journal of Ethnobiology and Ethnomedicine* 2013 **9**:23.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

